

Missing the point in noncommutative geometry

Nick Huggett^{a*}, Fedele Lizzi^{b,c,d†}, and Tushar Menon^{e‡}

June 23, 2020

^a Department of Philosophy, University of Illinois at Chicago
Chicago, IL, United States

^b Dipartimento di Fisica “Ettore Pancini”, Università di Napoli *Federico II*, Napoli, Italy

^c INFN, Sezione di Napoli, Italy

^d Departament de Física Quàntica i Astrofísica and Institut de Ciències del Cosmos (ICCUB),
Universitat de Barcelona, Barcelona, Spain

^e Faculty of Philosophy, University of Cambridge, Cambridge, United Kingdom

Abstract

Noncommutative geometries generalize standard smooth geometries, parametrizing the noncommutativity of dimensions with a fundamental quantity with the dimensions of area. The question arises then of whether the concept of a region smaller than the scale - and ultimately the concept of a point - makes sense in such a theory. We argue that it does not, in two interrelated ways. In the context of Connes’ spectral triple approach, we show that arbitrarily small regions are not definable in the formal sense. While in the scalar field Moyal-Weyl approach, we show that they cannot be given an operational definition. We conclude that points do not exist in such geometries. We therefore investigate (a) the metaphysics of such a geometry, and (b) how the appearance of smooth manifold might be recovered as an approximation to a fundamental noncommutative geometry.

*huggett@uic.edu

†fedele.lizzi@na.infn.it

‡tv288@cam.ac.uk

Contents

1	Introduction	2
2	Spectral triples	4
2.1	Meaning and definition	5
2.2	The spectral approach	7
3	Operationalism	14
3.1	The tempered operationalist approach to points	15
3.1.1	Classical space	16
3.1.2	Quantum kinematics	17
3.2	Noncommutative space	18
4	Ontology	20
5	Recovering space from NCG	24
5.1	The Moyal and Weyl representations	25
5.2	Finding Spacetime	28
5.3	Empirical coherence and physical salience	32
6	Conclusion	33

1 Introduction

In the vast landscape of contemporary theoretical physics, few research programmes come as close to engaging, as a matter of course, with traditional metaphysics as quantum gravity research programmes do. The two theoretical edifices that these programmes aim to unify (or replace)—quantum theory and relativistic gravitation theory—have been notoriously uncooperative with attempts at unification. It is not clear precisely which aspect of these theories is to blame, and what aspects ought to be held onto in future theories. Consequently, we are led back to traditional questions in the metaphysics of space and time. Questions like: what is the nature of space(time)? Are space(time) points fundamental? Is space(time) discrete?

In this paper, we discuss a particular approach to the metaphysics of discrete space suggested by one popular family of approaches to quantum geometry that go under the name of *noncommutative geometries*. From a philosophical perspective, attention to noncommutative field theories is valuable, because these theories allow us to embed our extant, well-confirmed physical theories in a broader logical landscape. Doing so allows us to unearth a number of tacit assumptions in

our current physical theories that might otherwise have been invisible, or appeared as matters of necessity.

Our goal, therefore, is to introduce noncommutative geometry to a wider philosophical audience, by discussing three metaphysical puzzles about the nature of space and, in particular, indeterminacy of location to which these geometries give rise. We understand ‘indeterminacy of location’ as referring to situations in which, for whatever reason, nature does not ascribe to a body a determinate matter of fact about its spatial location below a particular scale. The first puzzle, accordingly, is to characterise this particular brand of metaphysical indeterminacy. This leads to the second puzzle of how one ought to think about the ontology of a theory that is based on a noncommutative geometry. The final puzzle is to account for our experience of spacetime as, at least approximately, being described by a commutative geometry.

The family of approaches to modelling indeterminacy in quantum mechanics mentioned above broadly assert that if quantum mechanics is true, then particular facts about the world are ‘unsettled’—we can pose questions about the values of certain properties of systems such as, say, the x and the y components of spin, but nature itself does not determine the answers to such questions. Here, we focus on the subspecies of these approaches dubbed ‘supervaluationist’ (see e.g. [10, 11]).¹ We refer to this approach as modelling *indeterminacy as underdetermination*: one considers various precisifications of the models of the physical system, and models indeterminacy as the underdetermination of which of the precisifications truly represents the world.

Some sorts of quantum indeterminacy can plausibly be modelled as underdetermination, because ordinary quantum mechanics presupposes a continuous manifold of spatial points, so each of the precisifications is itself antecedently meaningful. Noncommutative geometry, on our interpretation, does not have the resources to make meaningful claims about localisability below a certain magnitude. We therefore argue that the indeterminacy that results from a noncommutative approach to spatial geometry, as suggested by noncommutative geometric approaches to quantum geometry is of a different kind. We call this *indeterminacy as meaninglessness*. In this paper, we cash out ‘meaninglessness’ in two distinct ways, depending on the approach to NCG: (i) §2 presents Alain Connes’ spectral triple generalisation of Riemannian geometry, and characterises meaninglessness as undefinability; (ii) §3 presents a concrete representation of quantum theories in noncommutative space, and characterises meaninglessness as non-operationalisability.

Having established our argument for a fundamental metaphysics that eschews the concept of arbitrary localisability, we discuss some alternative views of the ontology of a noncommutative field theory in §4. One thing all of these proposals agree on is that the elements of the relevant noncommutative algebra should be treated as fundamental. The picture of a field as an ascription of properties to points (supplemented with some story, kinematical or dynamical, about how

¹Since the ‘determinables-based’ approaches (primarily associated with Wilson and collaborators, [36, 7]) also presuppose a topological manifold structure on the domain of discourse, the criticism we offer in this paper also targets the determinables-based approach. However, for dialectical clarity, we choose to focus only on the supervaluationist approaches.

those points are related to each other) is untenable. While fields-first proposals have been in the philosophical literature for decades (Earman discussed so-called Leibniz algebras at least as far back as 1977 [16]), they have been presented as *alternatives* to standard ontologies for commutative theories like general relativity. In noncommutative field theories, fields-first interpretations are the only game in town.

In §5, we go on to examine a proposal for the recovery, from a noncommutative underlying geometry, of physical spacetime that is at least approximately commutative. In particular, we discuss a proposal that relies on structural features of quantum field theories to allow us, at least in some restricted but nonetheless physically salient circumstances, to recover a geometry that is approximately Minkowskian. The mismatch between the manifest and scientific images of space is an especially acute problem in the case of a theory with a putatively non-spatiotemporal fundamental ontology. This is because given that all evidence for any theory is ultimately in terms of spatiotemporal data. In recovering the manifest image from NCG, at least in a restricted context, we counter the possibility that NCG is *empirically incoherent*, to use a term introduced by Barrett [2] to describe theories whose truth undermines our justification for believing in their truth.

2 Spectral triples

A standard move in contemporary philosophy of spacetime is to model a spacetime theory as consisting of a smooth (i.e. infinitely differentiable), second countable, Hausdorff manifold on which are defined some tensor fields which encode spatial and temporal relations (in relativity theory, a Lorentzian metric tensor field) and some other tensor (or spinorial tensor) fields representing a matter distribution. The elements of the smooth manifold are typically treated as constituting the domain of discourse, call it M , of the theory; these elements, (commonly referred to as the ‘spacetime points’, although we choose not to use that term in what follows), are considered to be part of the fundamental, non-derived, non-emergent ontology of the theory.

The inclusion of particular mathematical structure in these models is justified, if only implicitly, by the idea that such structure allows us to represent structure that, plausibly, is part of the physical world. The metric tensor, for example, expresses our commitment to spatiotemporal lengths being relational properties of parts of the world. The topological structure in these theories plays a number of distinct roles, one of which is that it allows us to express a notion of localisation. In particular, we can speak of bodies as being localisable to an arbitrarily small interval in a topological manifold by considering smaller and smaller open sets, and claiming that the body is wholly located in each of them.

Even if we had reason to believe that it was not possible to localise a body to an arbitrarily small interval, call this the *localisability thesis*, it might still be the case that topological structure is needed to express that claim. On intuitions developed in the context of differential geometry, this might seem like a reasonable view: we talk of distances, whether finite or infinite, as being built out

of integrals of infinitesimal distances. And, indeed, this is how we set up the kinematics of ordinary quantum mechanics and quantum field theories: these theories still assume a fixed background spacetime with a particular topological (and a particular metric) structure. Claims about the impossibility of arbitrary localisations are ultimately cashed out in terms of the impossibility of localising to arbitrarily small intervals of the topological manifold, rather than in terms of some structure that dispenses with topological manifolds altogether.

Not only do we have good reason to doubt the localisability thesis, we believe that, even if it were true, the best way to cash out the localisability thesis, in general, would not be in terms of topological structure at all. In this section, we spell out our argument for this claim.

2.1 Meaning and definition

In order for an object, or a part of an object, to have a spatial location, it needs to be localised to some spatial region. An object is localised within some finite interval of length, call it δ (from which one can straightforwardly define the more useful notion of localisation to a finite area) iff the coordinate functions, x^i associated with the boundaries of that object satisfy the following constraint:

$$\delta \geq |x^i(p) - x^i(q)| \geq 0 \tag{1}$$

where $|\cdot|$ is a given norm in \mathbb{R}^n , $\delta < \infty$ and $p, q \in M$ are its boundary points.

The domain thus has metric structure, which is to say, enough structure to allow us to define a distance between any two of its elements. This conception of structure—a point set together with a privileged collection of coordinatisations—is familiar from standard practice in mathematical physics. It is worth, however, noting the subtle ways in which such structure is encoded in the formalism of first-order logic. That is the primary purpose of this subsection.

We begin by looking at how the story about this structure might be told in a mathematical textbook. As we noted, we can define a property like location because the coordinate functions are isometries from the domain, M , to \mathbb{R} . And when we specify a set of isometries, we pick out a particular equivalence class of metric spaces, and so our preferred coordinatisations can be thought of as inducing metric structure on M . And the something similar is true of topological, smooth, linear or any other form of geometric or algebraic structure. Let us call such a space a *structured domain*, and denote it as M_s . This induced metric structure on the structured domain allows us to talk of separations between locations in M_s in terms of separations of the images in \mathbb{R} of coordinate functions on points in M_s .

In terms of the formalism of first-order predicate logic, the story is different.² Here, we start with what we refer to as a *picture*, \mathfrak{A} .³ A picture for a first order language is pair $\langle M, \mathcal{I} \rangle$ where M

²For dialectical clarity, we discuss only first-order predicate logic in this paper. However, the most important idea introduced in this section, *definability*, along with the subsequent lemma about automorphisms, applies just as well to all higher-order logics. For more on this, see e.g. [19].

³What we call a picture, following Dewar [14], is more standardly referred to as a structure in logic texts.

2.1 Meaning and definition

is a domain of discourse and \mathcal{I} is an interpretation function. The domain of discourse is just a point set; it has no further structure, so we refer to this domain as the *bare domain*. The domain of the interpretation function is the set of nonlogical constants, a, a_1, a_2, \dots , n -place predicate symbols P, P_1, P_2, \dots and n -place function symbols, f, f_1, f_2, \dots . This set is known as the signature, call it Σ . The interpretation function is then:

1. Maps each n -place predicate symbol P to an n -ary relation $P^{\mathfrak{A}} \subseteq M^n$
2. Maps each n -place function symbol f to an n -ary operation $f^{\mathfrak{A}} : M^n \rightarrow M$
3. Maps each constant, a to an element $a^{\mathfrak{A}} \in M$.

A first order predicate calculus also includes variables, x, x_1, x_2, \dots —call this set V . A variable assignment α , is a function from the space of variables V , to the bare domain M .

In our specification of a structured domain, M_s , we relied on privileging certain coordinate functions. We could then characterise this structure by specifying its *automorphisms*—maps from M_s to itself that preserved the structure induced by the coordinatisation. We do something similar when characterising the structure using pictures, except that now, we consider automorphisms of a picture. Indeed, there is a bijection between automorphisms of pictures and automorphisms of structured domains, so, in what follows, we shift between speaking about pictures and structured domains.

Consider two pictures, $\mathfrak{A} = \langle D, \mathcal{I} \rangle$ and $\mathfrak{A}' = \langle D', \mathcal{I}' \rangle$, and a function $h : D \rightarrow D'$. Call such a function a *homomorphism* of the picture \mathfrak{A} iff for each n -tuple of D

- (1) for each n -place predicate $P: \langle a_1, a_2 \dots a_n \rangle \in P^{\mathfrak{A}}$ iff $\langle h(a_1), h(a_2) \dots h(a_n) \rangle \in P^{\mathfrak{A}'}$.
- (2) for each n -place function $f: h(f^{\mathfrak{A}}(a_1, a_2 \dots a_n)) = f^{\mathfrak{A}'}(h(a_1), h(a_2) \dots h(a_n))$.

If $\mathfrak{A} = \mathfrak{A}'$, then h is called an *endomorphism*. A bijective endomorphism is called an *automorphism*. We can thus specify a structured domain, M_s using a picture, \mathfrak{A} . Since it is the picture that specifies mathematical structure, we need to remember that certain useful concepts, like definability, are relative to a particular picture.

We can now (finally) speak precisely of what it is to be able to define an n -place relation (or function) *in a picture*:

An n -place relation R (on a set S) is **definable in a picture** if and only if the set of n -tuples of elements of the bare domain picked out by an interpretation function is invariant under the automorphisms of the picture, $\mathfrak{A} = \langle D, \mathcal{I} \rangle$.

However, this use risks confusion with the more general notion of mathematical structure, which we also discuss in the paper.

2.2 The spectral approach

The example below emphasises the role that the structure of the domain of discourse is playing here. For a finite domain, we can specify a relation by simply listing out n -tuples of elements of the domain. We cannot do that for domains of uncountable cardinality, as we have here. So we need to appeal to some further structure on the domain that will allow us to define a relation. Consider a manifold in which the only metric structure is induced via a ‘discrete coordinatisation’ of $N \subset M_s$, i.e. a map from a countable proper subset of M_s to the integers, \mathbb{Z} such that there is a smallest distance between any two points in N , denoted by μ . Any transformation on M_s which preserves the distances between the elements of N that are images of the discrete coordinate function is in this case an automorphism of M_s . Now consider the following putative relation on M_s , call it $R_\mu: \{\langle x, y \rangle \mid d(x, y) < \mu\}$. One characteristic feature of this relation is that if one element of a pair that constitutes R_μ is contained in N , then the other is not. Thus this relation is not invariant under all automorphisms of M_s : for example, an automorphism on M_s that leaves N unchanged, but permutes every other element of M_s . In this case, R_μ is not definable on the set N in a picture with domain M_s , as would need to be the case if we modelled indeterminacy of location as mere underdetermination of arbitrarily precise location.

Our restriction to a domain with this privileged discrete structure was contrived. But it was contrived in order to make an important point about representation and indeterminacy; the examples we discuss in the rest of the paper will be more physically motivated. Quantum indeterminacy in position of a particle on the de Broglie-Bohm interpretation, for example, can be unproblematically represented by an underdetermination between different models, where each model has associated with it a perfectly determinate notion of location, derived from the definability (in the technical sense) of arbitrarily short lengths (more precisely, a family of two-place relations on M corresponding to arbitrarily small separations). And this is perfectly permissible (indeed, encouraged) in a formalism which treats the structured domain as a metric space. In what follows, however, we will argue that a quantum mechanical approach to geometry mandates a noncommutative, discrete structure on the domain of discourse. As a result, separations below certain scales (again, more precisely, a family of 2-place relations on M corresponding to separations below a certain scale) will not be definable, so indeterminacy in position cannot be modelled by underdetermination between models with determinate notions of location.

2.2 The spectral approach

It appears, therefore, that we have a problem. It is all very well to say that a metric space can be characterised by assigning a real number to each pair of its elements, but how does one do this systematically for a domain M with uncountably many objects? We cannot simply list numbers associated with all pairs of elements in M . The standard move is to define a *line element* on M , $ds^2 = g_{ij}dx^i dx^j$, which represents, roughly, an infinitesimal displacement, and then integrate this line element along an arc that connects any two points in M in order to assign to that pair some real number. This ‘arc-connectedness’ is the basis of all differential geometry, and is an extremely

2.2 The spectral approach

powerful piece of mathematical technology.

Unfortunately, this move brings with it a problem: this definition of length commits us to being able to define lengths only if we can define arbitrarily short lengths, i.e. if the space is arc-connected. This won't do—recall from the previous section that we had good reasons to believe that lengths *above* a certain minimal scale ought to be definable even if lengths were not definable below that scale. In other words, we need to be able to define *some* lengths in a space that is not arc-connected, so our definition of length cannot require that a space be arc-connected. How ought one proceed in light of this demand?

Alain Connes' formalism of spectral triples promises to solve this problem. In order to assess the plausibility of this claim, we need to understand both the motivations and the mechanics behind this proposal. We begin by making precise the questions to which the spectral triple formalism provides the answer:

Question 1: What is the minimal structure of the domain of discourse required to make sense of locations?

We argued in the previous section that one natural answer to this questions is 'the structure of a metric space, $\langle M, d \rangle$ ', where d is the geodesic distance between any two elements in the domain of discourse. If the only way to turn a domain of discourse of uncountable cardinality into a metric space were by using the calculus of line elements, this answer would have been equivalent to 'the structure of a metric manifold, $\langle M, g_{ij} \rangle$, for domains of uncountable cardinality.' Our aim in this section is to build up a generalisable alternative characterisation of a metric space, and to answer in the affirmative the following question:

Question 2: Can we turn a domain of discourse of uncountable cardinality into a metric space even if it is not assumed to be arc-connected?

The goal, then, is to recover a metric space $\langle M, d \rangle$ algebraically, from a starting point that does not assume any metric or topological structure on the structured domain of discourse. Connes suggests using *spectral triples*, $\langle \mathcal{A}, \mathcal{H}, D \rangle$, where \mathcal{A} is a particular kind of algebra over a field, known as a *C^* -algebra*⁴, \mathcal{H} is a Hilbert space and D is a particular self-adjoint operator over that Hilbert space, known as a *Dirac operator*. The idea is simple: recast the structure that we know and love from differential geometry as special cases of these triples, check that we can still do differential geometry after this step, and then modify and deform these structures in such a way as to generalise differential geometry to unfamiliar domains.

We split the task of recasting differential geometry in the language of spectral triples into two steps:

⁴A C^* -algebra is an algebra over a complex field, equipped with an *inner involution*, denoted by \star , which generalises the operation of taking the complex conjugate of a number, and a *norm*, which generalizes the modulus of the complex numbers, with respect to which the algebra is complete. Complex number themselves are the simplest example of C^* -algebra.

2.2 The spectral approach

Step 1: Recover a topological manifold, M , from a spectral triple.

Step 2: Recover a geodesic distance function, d on M , from a spectral triple.

Step 1 is completely straightforward and it builds on mathematical work in the forties, mainly a well-known representation theorem due to Gelfand and Naimark [23], which we describe briefly. Consider a Hausdorff topological space, for simplicity we may think of a manifold, M , for instance the (2-dimensional) plane or sphere. Defined on it are the scalar fields, continuous functions that assign a complex number to each point: the set of such functions⁵ is known as $C(M)$. To understand the following, it is important to distinguish between such fields, which are functions over all space, from their values at a point: the former are complete ‘configurations’ of individual point-values.⁶

Two scalar fields, ϕ and $\psi \in C(M)$, can be multiplied together in an obvious way to obtain a third, $\chi \in C(M)$ – the value of χ at any point p , is just the ordinary product of the values of ϕ and ψ at p : $\chi(p) = \phi(p) \cdot \psi(p)$. Such ‘pointwise’ multiplication of fields is in fact so obvious as to almost be invisible: how could there be an alternative? In §5.1, we shall see that there are alternative rules for multiplying fields, and indeed, they may even be more physical than pointwise multiplication.

Because ordinary multiplication is commutative, $a \cdot b = b \cdot a$, so is pointwise multiplication for elements of $C(M)$: $\phi \cdot \psi = \psi \cdot \phi$. Nonetheless, the algebra contains a great deal of information about the space on which the fields live. In fact, the algebra contains *all* the information that we typically take to characterize a topological space. Topology, understood as characterizing relations between points, can be reconstructed from purely algebraic data as maximal ideals⁷, the neighborhood of a point can be likewise inferred from the relations among ideals. Global characteristics are also encoded in the algebra; for example a closed compact space (such as a circle) is described by an algebra which contains a multiplicative identity element. By contrast open spaces such as the real line are described by algebras which lack such an element. In short, this representation theorem states the logical equivalence of a space topology and its algebra $C(M)$.

It’s worth emphasizing the strength of this point, by reflecting on what is meant by an ‘algebra’: nothing but a pattern of relations – a structure – with respect to some abstract operations. One might, for instance fully characterize an algebra by saying that there are two elements, $\{a, b\}$, and an operation \circ , such that $a \circ a = b \circ b = b$ and $a \circ b = b \circ a = a$ (and specifying that the operation is associative). What the elements are is not relevant, neither is the meaning of \circ ; all

⁵For our purposes we need not go into detail as for the class of functions, continuous (for topology) or smooth (for differentiability) or other classes.

⁶The usual notation for fields, e.g. $\Psi(x)$, tempts a conflation here, as the argument could be read as a particular value; but it instead indicates that we have a function over (co-ordinatized) points. Since that is understood, we might leave out the argument, and just denote the field (the function, the configuration) as Ψ . With respect to the set C^∞ , the fields are its elements, and their identities depend on the point-values (two fields are the same field, iff all their point values agree).

⁷An ideal is a subalgebra such that the product of one of its elements by any element of the algebra still belongs to the ideal.

2.2 The spectral approach

that matters is how many elements, and what function on pairs of elements \circ is. Of course, an algebra can have different concrete *representations*: concretely, a might be represented by the set of true propositions, and b by the set of false propositions, in which case \circ is represented by the boolean not-biconditional connective. But there are other representations: addition mod-2 for instance (and perhaps a could be represented by the presence of a 30kg hemisphere of uranium 235, b by the absence, and \circ by the operation of putting together – the critical mass of U^{235} is 52kg!). These are different representations of a *single* algebra, which captures their common structure.

It is not relevant that the concrete elements of the algebra are fields over the manifold, all that need be specified are their relations with respect to a binary operation. However, the scalar fields on a particular manifold define a specific algebra, and, according to the representation theorem, no other manifold has scalar fields with the same algebra. The point is that the algebra does all the work: there is nothing smuggled in about the manifold simply because we realize the algebra with fields over it. Additionally, every abstract commutative C^* -algebra can be represented as an algebra of continuous functions $C(M)$ over some Hausdorff space M . So that settles our choice of algebra in the spectral triple: let \mathcal{A} be $C(M)$, the algebra of continuous complex-valued functions over M . In general, \mathcal{A} determines a Hilbert space as well via a procedure known as the *GNS construction*.⁸

Step 2 is a little more involved, and it follows step 1 by nearly half a century, thanks to the works of Alain Connes and others. It requires some ingenuity. As we noted above, we cannot immediately recover, from the algebra of continuous functions, a metric manifold $\langle M, d \rangle$ in the way that we could recover a compact topological manifold M . One might wonder, however, if there is some subalgebra of $C(M)$ that might encode metric facts about M . And there is, but this algebra can only be picked out if we allow ourselves the third piece of structure in a spectral triple, the *Dirac operator*, D . If we stipulate that our domain of discourse, $\langle M, d \rangle$, has enough structure for us to define a notion of spinors, roughly speaking, we can then define a differential operator, $D := i\gamma^\mu \partial_\mu$ on the vector space of these spinors.⁹ With this extra structure, we have enough machinery to isolate a subalgebra of \mathcal{A} that will allow us to recover d , the geodesic distance.

Consider the subalgebra of $C(M)$ known as the (algebra of) *Lipschitz functions*, defined as follows:

Given two metric spaces, $\langle M, d_M \rangle$ and the real line \mathbb{R} , the function $f : M \rightarrow \mathbb{R}$ is a **real-valued Lipschitz function** if and only if there exists a real-valued constant, K such that:

$$|f(x_1) - f(x_2)| \leq K d_M(x_1, x_2)$$

⁸Briefly, this is achieved by treating the underlying vector space of \mathcal{A} as its own representation space and using the representation, together with a specification of a complex-valued functional, ω (more on this object in §3.2) to induce an inner product on the vector space. Finally, one takes care to ensure that the space is completed in the associated norm.

⁹Less roughly speaking, we define a differential operator on the Hilbert space of square integrable spinors.

2.2 The spectral approach

Since Lipschitz functions can only be defined on metric spaces, the idea is that if we can find a subalgebra of \mathcal{A} of Lipschitz functions, \mathcal{A}_L , we could use that algebra to reconstruct the geodesic distance on M . The problem thus splits into two parts:

Part 1: Identify the subalgebra of $C(M)$ that constitutes the algebra of Lipschitz functions that we denote as $C_L(M)$.

Part 2: Identify the appropriate Lipschitz function for every pair of points on M such that one can identify it with the geodesic distance between those two points.

For part 1, we start by considering a bounded measurable function, $a \in C(M)$, on M . From the GNS construction associated with this algebra, we know that it can be represented as an operator on a Hilbert space.¹⁰ There is a theorem that states that this function will be almost everywhere¹¹ equal to a Lipschitz function if and only if the operator $[D, a]$ is bounded [9], where D is the Dirac operator defined on the Hilbert space \mathcal{H} . So if we specify a Dirac operator, we can identify the algebra of Lipschitz functions, $C_L(M)$, a proper subalgebra of $C(M)$.

For part 2, we recall, from the definition of a Lipschitz function, that, for the value $K = 1$, the norm of the difference of image points is the geodesic distance. We are thus led to the following suggestion for the geodesic distance function on M :

$$d(p, q) = \sup\{|a(p) - a(q)|; a \in C(M), \|[D, a]\| \leq 1\} \quad (2)$$

where $|\cdot|$ is the L^2 -norm on \mathbb{C} and $\|\cdot\|$ is the norm on the Lipschitz algebra $C_L(M)$. Note that we are defining a distance function on M indirectly—by appealing to structure in the algebra $C_L(M)$. We find that this agrees with the standard definition of geodesic distance in terms of line elements in an arc-connected space:

$$d_M(p, q) = \inf_{\gamma} \int_{\gamma} g_{ij} dx^i dx^j \quad (3)$$

where γ is a path from p to q .

The advantage of the use of spectral triples, in the context of the discussion in the previous section, is clear—we can, using equation ((2)), define a notion of distance between two elements of a domain of discourse *even when that domain is not assumed to be arc-connected*. We thus have the construction that we required in order to answer question 2 in the affirmative. Crucially, since we no longer need to assume M is arc-connected, we can generalise the algebra, from the commutative $C(M)$ to a noncommutative \mathcal{A} . We do not need to worry that there is no longer a sensible notion of infinitesimal distance. All we need is a determinate specification of geodesic distance between elements of the subset of the domain of discourse *for which the notion of separation is definable*. This specification does not need to piggyback on a specification of infinitesimal

¹⁰To cut down on unnecessarily complicated notation, we refer to this element as a , while remaining vigilant about correctly identifying which space these similarly labelled operators belong to.

¹¹i.e. everywhere except for a measurable set of Lebesgue measure zero.

2.2 The spectral approach

distance, and is thus still available to us in spaces characterised by having a noncommutative algebra of functions.

At this stage, we do no (serious) harm by thinking of M as a set of possible locations of a single particle; we will have to generalise this shortly. From the discussion above, we know that $C(M)$ has the structure of a complex algebra, whose multiplication action is determined by pointwise complex-number multiplication. Consider the set $\mathfrak{S}(C(M))$ of nonzero C^* -algebra homomorphisms between $C(M)$ and \mathbb{C} . This set is known as the *Gelfand spectrum*, and consists of maps, ρ whose action is defined as follows:

$$\rho(a) = a(p) \in \mathbb{C} \forall p \in M$$

Elements of the Gelfand spectrum therefore stand in a one-one correspondence with points of the domain M of complex continuous functions. In fact this bijection preserves topological structure: the Gelfand spectrum is homeomorphic to the set M . Given this homeomorphism, one can find an expression for the distance function d_M on M in terms of elements of the Gelfand spectrum:

$$d_M(p, q) = d_{\mathfrak{S}(C(M))}(\rho, \eta) = \sup\{|\rho(a) - \eta(a)|; a \in C(M), \|[D, a]\| \leq 1\}$$

where $\rho, \eta \in \mathfrak{S}(C(M))$ are the elements mapped to p, q respectively by the homeomorphism between $\mathfrak{S}(C(M))$ and M . Note that the notation now makes clear that the elements of $C(M)$ are arguments of the elements of the Gelfand spectrum.

Denote the convex subsets of a generic algebra \mathcal{A} as $S(\mathcal{A})$; and the ‘extreme boundary’ of a convex set K , as $\partial_E(K)$.¹² Elements of the extreme boundary of $S(\mathcal{A})$ are also known as *pure states*. It turns out that the space of pure states is also homeomorphic to M when \mathcal{A} is commutative. So now we have three spaces, M , $\mathfrak{S}(C(M))$, and $\partial_E S(C(M))$, arrived at by different constructions, that are homeomorphic to each other.¹³ Let us focus on one important pair, $\partial_E S(C(M))$ and M . This isomorphism means that points in M stand in a one-one correspondence with pure states. Consider the appropriate expression for d on $\partial_E S(C(M))$:¹⁴

$$d_{\partial_E S(C(M))}(\alpha, \beta) = \sup\{|\alpha(a) - \beta(a)|; a \in \partial_E S(C(M)), \|[D, a]\| \leq 1\}$$

All of this demonstrates that we should not be fooled into thinking that a geodesic distance can only be defined when M and $\partial_E S(\mathcal{A})$ are homeomorphic—once this isomorphism is broken by replacing a commutative algebra with a noncommutative algebra of functions on M , $\partial_E S(\mathcal{A})$ is still a metric space with a metric given by equation (2). But now, this metric space is no longer

¹²A vector space K is called a convex set iff $\forall v, w \in K$ and $t \in [0, 1]$, $tv + (1-t)w \in K$. Its extreme boundary is the set of all $v \in K$ such that if $v = tw + (1-t)x$ for certain $w, x \in K$ and $t \in [0, 1]$, then $v = w = x$.

¹³We can add a fourth: the set of probability measures on M .

¹⁴It is worth noting that this expression gives us distances on the entire space of states on $S(C(M))$, not just the pure states. However, in this section, we focus just on the subset of pure states.

2.2 The spectral approach

isometric to the space $\langle M, d_M \rangle$ (unsurprisingly, given that they are no longer even homeomorphic). Consequently, and crucially, the pure states are no longer identified with points in M (the choice of terminology is not accidental—in quantum mechanics, these are the standard pure states that can be identified with rays in the system’s Hilbert space).

To reiterate, when $\mathcal{A} = C(M)$, which is commutative, the space of pure states, $\partial_E S(C(M))$ is homeomorphic (and can be made isometric) to the space M , so it does not matter which space we begin with. This is no longer true when \mathcal{A} is noncommutative. In this case, the geodesic distance function still maps pairs of pure states to real numbers, but the space of pure states, $\partial_E S(\mathcal{A})$ is no longer homeomorphic to the domain of the functions that constitute \mathcal{A} . The geodesic distance function now identifies distances between pure states which, in quantum mechanics for example, cannot be interpreted as points of space. This is what underlies the ‘pointlessness’ of NCG, as alluded to in several discussions of NCG, for example:

The concept of a point becomes evanescent, and in some cases one is forced to abandon it altogether. [31, p. 95]

The conceptual shift in NCG is to treat the algebra \mathcal{A} as fundamental, and the structure on the space X as derived (we will explore some of the metaphysical consequences of this move in §4). In regimes where we need only focus on commuting algebras of observables, distances between pure states can, to whatever the appropriate degree of approximation, be identified with distances between locations.

We are now in a position to understand how this construction allows us to determine under what circumstances the 2-place ‘separation’ relation is definable. Define an automorphism, h of a spectral triple as an automorphism of A such that leaves the Dirac operator invariant. If $\mathcal{A} = C(M)$, then D picks out the same distance function before and after the transformation. In other words, h is an automorphism of $\langle C(M), \mathcal{H}, D \rangle$ iff

$$d(p, q) = \sup\{|h(a(p)) - h(a(q))|; a \in \partial_E S(C(M)), \|[h(D), h(a)]\| \leq 1\} \quad (4)$$

where $d(p, q)$ is defined as in equation (2).

This $d(p, q)$ is equal to the distance defined directly on M by equation (3). Consider a relation on $\langle M, d_M \rangle$:

$$R_\mu := \{p, q \in M | d_X(p, q) = \mu\}$$

where $d_M(p, q) = \inf \int_\gamma g_{ij} dx^i dx^j = \mu$ and $\mu \in R$. This relation is invariant under an automorphism h on $\langle M, d_M \rangle$. This automorphism will induce an automorphism h_A on $\langle C(M), \mathcal{H}, D \rangle$ such that:

$$\sup\{|h_A(a(p)) - h_A(a(q))|; a \in \partial_E S(C(M)), \|[h_A(D), h_A(a)]\| \leq 1\} = \mu.$$

We can define a relation $R_\mu^{C(M)}$ on $\partial_E S(C(M))$. This relation is invariant under all and only the automorphisms that preserve R_μ . They are completely equivalent. So it does not matter whether

we privilege an algebraic or geometric description. They agree on the magnitudes that we are interested in here—lengths and areas, and consequently, as μ is made arbitrarily small, both $R_\mu^{C(M)}$ and R_μ pick out the same relation.

But we are interested in what happens when we leave these regimes and instead consider noncommutative algebras of observables. Let θ quantify the magnitude of the noncommutativity of \mathcal{A} , i.e. $\forall \hat{x}, \hat{y} \in \mathcal{A}, [\hat{x}, \hat{y}] = i\theta$. Let \mathcal{A} be noncommutative, and at the same time, let us structure M as a metric space, $\langle M, d_M \rangle$. In this case, as mentioned above, M will no longer be the space of pure states, $\partial_E S(\mathcal{A})$. We can, of course, still define the relation R_μ on $\langle M, d_M \rangle$ because it remains invariant under automorphisms, h of $\langle M, d_M \rangle$. But things change when we take the algebra \mathcal{A} as fundamental. In that case, the putatively equivalent relation $R_\mu^{\mathcal{A}}$ would have to be expressed in terms of the structures in $\partial_E S(\mathcal{A})$:

$$R_\mu^{\mathcal{A}} := \{ \phi, \psi \in \partial_E S(\mathcal{A}) \mid d_{\partial_E S(\mathcal{A})}(\phi, \psi) = \mu \}$$

where $d_{\partial_E S(\mathcal{A})}(a, b)$ is given by equation (2).

The problem is that while this relation is, indeed, invariant under automorphisms of the noncommutative spectral triple, $\langle \mathcal{A}, \mathcal{H}, D \rangle$, it is no longer equivalent to the relation R_μ . This is because although the metric space of pure states $\langle \partial_E S(\mathcal{A}), d_{\partial_E S(\mathcal{A})} \rangle$ is invariant under spectral triple automorphisms, it is no longer isometric to $\langle M, d_M \rangle$. So R_μ and $R_\mu^{\mathcal{A}}$ are invariant under different automorphisms: they are different relations. Moreover, unlike in the commutative case, $R_{m\mu}^{C(M)}$ is not invariant under the automorphisms of the picture that preserve R_μ . We can, however, restrict our attention either to commutative algebras, or to regimes in which the algebra of relevant observables can be treated as being commutative (i.e. the scale μ is much larger than the noncommutation factor θ). In these scenarios, $R_\mu^{\mathcal{A}}$ is once again equivalent to (or nearly equivalent to) R_μ . The upshot of this discussion is that R_μ , and hence our concept of localisation, is definable in a theory whose domain of discourse is a spectral triple only if the algebra is commutative.

3 Operationalism

Philosophers of physics with operationalist leanings might be sympathetic to but nonetheless unmoved by the argument in the previous section. Operationalism is a view about how words that describe concepts acquire meaning, according to which ‘the concept is synonymous with the corresponding set of operations.’ [4] From the perspective of a physical theory, rather than trying to imbue an abstract formalism with physical salience, it is built into the formalism from the beginning.

Operationalism was introduced by Percy Bridgman, and most famously discussed in his *The Logic of Modern Physics*. Although Bridgman himself does not present his view as such, it can be read as advancing a specific semantic claim about scientific words, namely that the meaning of a

3.1 The tempered operationalist approach to points

word is (in a sense to be made precise) completely exhausted by a specification of the operations that one would need to perform in order to measure a magnitude corresponding to that concept. Of particular relevance to the discussion in this paper is the following quote from Bridgman:

If a specific question has meaning, it must be possible to find operations by which an answer may be given to it. It will be found in many cases that the operations cannot exist, and the question therefore has no meaning. [4, p. 28]

For various reasons, as a semantic theory, operationalism is no longer fashionable amongst contemporary philosophers. However, the empiricist spirit of operationalism still underwrites a standard approach among physicists and philosophers of physics for clarifying obscure concepts. This approach often proceeds via some form of thought experiment: think of Newton's globes as a way of understanding absolute space or the behaviour of light rays and test particles as a way of understanding the chronogeometric significance of the metric in general relativity.

It is in this spirit that, in this section, we present what might best be termed a 'tempered operationalism', according to which a necessary condition on a concept having physical content is that it is possible, by the lights of physical theory, to describe a (perhaps idealised) measurement procedure for a magnitude associated with the concept. We refer to such concepts as **operationally definable**. To give operationalism substance, one has to specify what measuring operations are available; since we are interested in the possibility of operationalizing points of space, we will consider a (probabilistic) particle detector.

3.1 The tempered operationalist approach to points

Having clarified the sense in which we understand operationalism, in this section, we describe an idealised location measurement procedure. We argue that, on this setup, in ordinary classical field theory, localisations can be arbitrarily precise in principle, but trouble starts to brew in noncommutative geometries, leading to a violation of **operational definability**.

In brief, the argument as follows. In classical field theories on a commuting space, any uncertainty in measurement of location is down to technological or epistemic limitations. To account for these limitations, we associate what one might call an *epistemic state* with a particle: a probability measure over some interval of space representing our uncertainty about the exact location—the *ontic state*—of a particle. We then model an idealised measurement in such a way that the probability distribution that characterises our epistemic state becomes infinitely peaked at the particle location, in the limit that the uncertainty tends to zero.

In theories of noncommutative space, we again assume—but this time for *reductio*—that there is an ontic state corresponding to an arbitrarily precisely localised particle. We construct the analogue of an epistemic state: a density operator. We then attempt to localise this epistemic state to an arbitrarily small area and discover that this leads to ascriptions of negative probabilities. Since these measures are not elements of the state space, this signals a pathology. The only

3.1 The tempered operationalist approach to points

way to avoid this pathology, we argue, is to drop the assumption that there is an ontic state corresponding to an arbitrarily precisely located particle. Thus, even in principle, it is not possible, in a noncommutative space, to localise a particle below a certain area. Operationally, then, such areas—and *a fortiori* points—are undefinable.

3.1.1 Classical space

We will thus operationalize spatial regions and points in terms of particle location measurements; if measurements are only physically possible to finite precision, we argue that areas smaller than that scale do not exist. So first, consider a single particle prepared in a state localised to a region in M . In practice, there will be some uncertainty because of technological limitations in the preparation the system. Second, we model a location measuring device: some apparatus which “clicks” if the particle is in a region, and is silent otherwise. Again, in practice there will be uncertainty in the measurement, because of our lack of full control of the dynamics. Because of both kinds of uncertainty—and for technical and dialectical reasons—we model both the state of the particle, and the measuring device with Gaussian functions.

Consider, first, the commutative case. Here, the coordinate functions of the two dimensions commute and therefore decouple (we will shortly see why this is not the case in noncommutative space), so it is sufficient to consider just one dimension. We represent the state of a particle as a *normalised* Gaussian centered around a point x_0 :

$$\rho = \frac{1}{\sqrt{\pi\alpha}} e^{-\frac{(x-x_0)^2}{\alpha^2}}$$

where α is the width of the Gaussian.

Observables are represented in this formalism as real functions of configuration space, say $f(x)$, and a measurement will give as average value $\langle f \rangle_\rho = \int dx \rho(x)f(x)$. The limit $\alpha \rightarrow 0$ is well defined and $\lim_{\alpha \rightarrow 0} \langle f \rangle_\rho = f(x_0)$ For instance, for position, x :

$$\langle x \rangle_\rho = x_0 ; \langle x^2 \rangle_\rho = x_0^2 + \frac{\alpha^2}{2}$$

so that the uncertainty is $\Delta x = \frac{\alpha}{\sqrt{2}}$ The limit $\alpha \rightarrow 0$ is well defined, and in that limit, the state is perfectly localised in x_0 .

We also model a position measuring device, with resolution β as a normalised Gaussian.

$$\frac{1}{\sqrt{\pi}\beta} g_1(x) = e^{-\frac{(x-x_1)^2}{\beta^2}}$$

The average value of the corresponding observable, for $\alpha = \beta$ is then

$$\langle g_1 \rangle_\rho = \frac{\beta_1}{\sqrt{2\alpha^2}} e^{-\frac{(x_0-x_1)^2}{2\alpha^2}} \quad (5)$$

This quantity is very small unless $x_0 \sim x_1$, i.e. the device is near the particle. When α is sufficiently small this quantity is close to one, and goes to zero very fast as $x_0 - x_1$ grows. In other words, this observable indeed corresponds to a position measurement, capable of discriminating the location of a particle to arbitrary accuracy, as $\alpha \rightarrow 0$. We will see later (§5.1) that things are quite different in noncommutative geometry.

3.1.2 Quantum kinematics

In the previous section, we were interested in the classical kinematics of location. Consequently, our states were just elements of some configuration space, observables were functions of these states, and measurement outcomes (magnitudes) were elements in the range of these functions. Quantum kinematics is different. The textbook story¹⁵ is that states are positive trace-class operators¹⁶ on a Hilbert space, observables are self-adjoint operators on that Hilbert space and measurement outcomes are probabilities over magnitudes and are determined by the Born Rule. In particular, the expectation value associated with some observable, call it \hat{O} is:

$$\langle \hat{O} \rangle_\rho = \text{Tr}(\rho \hat{O})$$

where ρ is a state.

If we assume that negative probabilities signal a pathology (or contradiction) in our theory, we need to strengthen the physicality condition on the state. It is not sufficient that ρ be merely positive trace class. One needs to add the further constraint that it have no negative eigenvalues. Thus, we stipulate a condition of **state physicality**:

State Physicality: Physical states of a quantum system are: (i) positive trace class operators on complex Hilbert spaces and (ii) have no negative eigenvalues.

If we begin with a set of self-adjoint operators on a complex Hilbert space, we can define an operator norm on this set, and endow this set with the structure of a C^* -algebra. We can incorporate some of the structure of the Born Rule into the definition of a state by defining states as maps from the C^* -algebra of operators into the field of complex numbers. More precisely, given an algebra with a norm, a *state* ρ is a map from the elements of the algebra into the set of

¹⁵as presented in e.g. [33].

¹⁶A state represented by an operator in quantum mechanics is trace class if it has a well-defined (i.e. basis-independent) trace. If this trace is positive, then the operator is positive trace class.

complex numbers with the following properties:

$$\begin{aligned}\rho(\alpha_1 f_1 + \alpha_2 f_2) &= \alpha_1 \rho(f_1) + \alpha_2 \rho(f_2) \\ \rho(1) &= 1 \\ \rho(f^+ f) &\geq 0 \\ \|\rho\| \equiv \sup_{\|f\| \leq 1} \rho(f^+ f) &= 1\end{aligned}$$

C^* -algebras allow us to express the kinematics of classical and quantum mechanics in broadly analogous ways. We should flag, however, the contentious nature of the appellation ‘quantum’. For some, e.g. [29], it is sufficient for a theory to be quantum that the algebra of observables be noncommutative. Others [20], argue that the quantum/classical divide is less clearly delineated, and lies on a parametrised spectrum, where the actual divide is sensitive to other considerations. The cases that we consider in this section are noncommutative, but are not ‘quantum’ in the strict sense of imposing \hbar -dependent commutation relations on the canonical variables. In this paper, we remain neutral on the substantive question, but for clarity speak only of commutative and noncommutative field theories, reserving our use of ‘quantum’ for later (§5.2), for domains that can strictly be thought of as quantum mechanical. But the point here is that the framework just developed is appropriate for any situation with noncommutative observables, and so is appropriate to the case of noncommutative *space*, as we shall now see.

3.2 Noncommutative space

With the algebraic characterisation of geometry in mind, we now model the measurement procedure as we did in §3.1.1, *mutatis mutandis*, for a noncommutative space. We do not need the full structure of the spectral approach described in §2.2, instead we work with a simple special case. We should stress at this stage that the lessons that we draw about measurements in this simplified model, based as they are on claims about the noncommutativity of the algebra of coordinates, generalise to more other systems modelled by noncommutative spectral triples.

Consider a simple toy example of a space in which what were previously coordinate functions (what we henceforth refer to as ‘base elements’) are now self-adjoint operators with the commutator:

$$[\hat{x}, \hat{y}] = i\theta.$$

As before, the area θ is the measure of the noncommutativity, analogous to \hbar , in the canonical commutation relations. \hat{x} and \hat{y} are of course fields, and of this relation, the algebra of *functions* of \hat{x} and \hat{y} —the field algebra \mathcal{A} —which in the classical case was commutative, is now noncommutative. In particular, their polynomials are noncommutative elements of the algebra.

So we seek a representation of the algebra, satisfying the quantum kinematics described above, via a ‘quantization map’ that associates to any function an operator. There is some freedom in

3.2 Noncommutative space

this choice, and usually one uses the *Weyl map*, which associates self-adjoint operators to real functions. The definition is made via the Fourier transform

$$\tilde{f}(\xi, \eta) = \int dx dy f(x, y) e^{\frac{i}{\theta}(\xi x + \eta y)}$$

the operator corresponding to the function $f(x, y)$ is therefore

$$\hat{F} = \int d\xi d\eta \tilde{f}(\xi, \eta) e^{-\frac{i}{\theta}(\xi \hat{x} + \eta \hat{y})}.$$

The inverse map, which associates functions to operators, is called the *Wigner map*:

$$\tilde{f}(\xi, \eta) = \text{Tr} F e^{\frac{i}{\theta}(\xi \hat{x} + \eta \hat{y})} \quad (6)$$

The operators we obtain can always be represented as acting on a separable Hilbert space, so both operators and states are represented as (infinite dimensional) *matrices*. Then the requirements (6) translate to the claim that we can associate with a state an Hermitean matrix $\hat{\rho}$ with positive eigenvalues and $\text{Tr} \hat{\rho} = 1$. This is standardly referred to as a density matrix.

Consider now a classical Gaussian epistemic state peaked around the origin and of width α :

$$\rho = \frac{1}{\alpha^2 \pi} e^{-\frac{x^2 + y^2}{\alpha^2}}$$

The quantization map associates an operator to this function, and we may write this operator in the x basis as:¹⁷

$$\hat{\rho} = \frac{1}{\alpha^2 \pi} e^{-\frac{x^2 + \theta^2 \partial_x^2}{\alpha^2}}$$

This operator has well defined action on functions of x .

This matrix has been calculated ([13, 30], and found to be:

$$\hat{\rho} = \begin{pmatrix} \frac{2\theta}{\alpha^2 + \theta} & 0 & 0 & \dots \\ 0 & \frac{2\theta}{\alpha^2 + \theta} \left(\frac{\alpha^2 - \theta}{\alpha^2 + \theta} \right) & 0 & \dots \\ 0 & 0 & \frac{2\theta}{\alpha^2 + \theta} \left(\frac{\alpha^2 - \theta}{\alpha^2 + \theta} \right)^2 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

This operator is trace class, with $\text{Tr} \hat{\rho} = 1$ in all cases. So it satisfies the first condition of state physicality. But under certain circumstances, it fails the second: for certain values of α , it gives rise to negative expectation values for any observable. This can be inferred from the eigenvalues of this operator, for particular values of α and θ .

$\alpha > \sqrt{\theta}$ The state is not particularly localised, and $\hat{\rho}$ is a density matrix, with all eigenvalues positive

¹⁷Remember that x and y do not commute, and therefore do not have a simultaneous basis, hence we cannot write the operator using both of them.

and smaller than 1.

$\alpha = \sqrt{\theta}$ The state is a density matrix with the first eigenvalue equal to 1, and the rest vanishing.

$\alpha < \sqrt{\theta}$ The matrix *is not* a density matrix, the first eigenvalues is larger than 1, and negative eigenvalues appear, which lead to negative expectation values for all operators. Thus the operator violates **state physicality**.

Thus the quantum state corresponding to a classical epistemic state localised to an area below $\sqrt{\theta}$ does not correspond an ontic quantum state—the associated density operator does not represent a physical state.

To better understand this result, consider the position measurements that we take to operationalize position in the noncommutative framework. $\langle \hat{x} \rangle = 0$, which reveals no problem, but if we consider higher powers we find:

$$\langle \hat{x}^4 \rangle = \frac{3}{16}(2\alpha^4 - \alpha^2\theta - \theta^2) \quad (7)$$

which—impossibly—is negative for $\alpha < \sqrt{\theta}$. Moreover, that the value of $\langle \hat{x} \rangle$ is positive is in fact an artifact of the symmetric state we chose; a different ρ , not symmetric around the origin, would have shown pathological behaviour even for a direct position measurement. In §5.1 we will introduce the formalism necessary to also see the noncommutative analogue of 5, a position measurement involving an imprecise device. We will see in more concrete terms why we cannot operationalize position measurements: attempts to measure below a certain scale are frustrated.

We conclude that measurements attempting to localize particles to a linear scale smaller than $\sqrt{\theta}$ are unphysical; of course, this result is a reflection of the fact that there are no ontic states localizing particles below this scale. Then, from our tempered operationalism, we conclude that space in fact has no regions smaller than this scale – and in particular, is ‘pointless’.

4 Ontology

We have argued that the notions of point, or region smaller than the commutation scale are undefinable in noncommutative geometry; undefinable in the formal sense of definition in a picture, and undefinable in the physical sense of tempered operationalism. All we have is a (metric) space of (pure) states, and these can only be identified as being pointlike if the algebra of fields is commutative. In that case, the traditional conception of the manifold fails too, and with it the ‘fields as properties of spatiotemporal points’ view about the ontology of a field theory (as endorsed by, e.g. Hartry Field [21]). Thus the question of the interpretation of theories of noncommutative geometry arises: what kind of world, what kind of basic elements, does it describe? One immediate thought is to develop a structure of spacetime by composing *regions* each of which is of size above the commutation scale. The issue with this proposal is that the set theoretic closure of such open

sets includes regions smaller than the commutation scale; so such an interpretation would require non-standard laws of composition. We take such issues as indicating that such an approach would shoehorn classical notions into what is, essentially, a quantum theory. So we will instead propose that the algebra, \mathcal{A} , whose elements are not intrinsically spatiotemporal at all, is itself the basic ontology. Naturally, this will require some elaboration!

Note that until now, we have only spoken of spatial, rather than spatiotemporal noncommutativity. From this point on, we also indulge in talk of noncommutative ‘spacetime’. But note that we always take the time coordinate to commute with the spatial coordinates (even when the latter do not commute with each other).¹⁸ We will also refer to noncommutative geometry as ‘NCG’ for brevity.

In Book IV, Chapter 1 of his *Physics* Aristotle offers the view that existence requires being somewhere: everything that is, has a place. (He is setting up the question, attributed to Zeno, of where places are, if they exist.) This idea is intuitive: the world seems fundamentally spatial, and it starts to capture the idea that ‘real’ things can be interacted with, by traveling to them. If one accepts such a view of existence, then it becomes impossible to take the algebraic formulation of the theory as giving an ontology for the world, because it does away with space as a fundamental object. Modern philosophers are more likely to take a logical view of existence, perhaps adopting Quine’s view that to be is just to be the value of bound variable in a true theory. At any rate, seems inadvisable to reject the algebraic formulation out of hand because of a view of existence.

Related to the idea that ties existence to space, is the idea that comprehension requires the spatial. Maudlin [32], following Barrett [2], expressed a concern that theories without fundamental spatiotemporal quantities could not be connected to experiments, which immediately concern local beables. But a vaguer objection that only spatiotemporal theories can be properly ‘understood’ perhaps remains; useful predictions might be possible, but otherwise a theory can only be an instrument, not comprehensible by us. Kant had a view like this of course, which influenced Maxwell in the construction of electromagnetism; and such claims were made by Schrödinger in his arguments with Heisenberg. See ([12]) for some history of the topic. But such instrumentalism is no more of a threat to an algebraic interpretation than Aristotle’s spatial view of ontology. First, one can make the same kind of move to logic as before. Suppose a theory is algebraic rather than geometric, then it may not give an image that is easily visualized by the human mind, but it still can provide understanding in the sense of systematizing the connections between different parts of nature; between the quantum and gravitational realms, ideally. That is, our ability to understand formal systems that aren’t spatial does plausibly give us the ability to understand physical systems that are not spatial. Second, there is a sense of ‘understanding’ that indicates facility with a theory rather than any sense that its models can be visualised. For instance, de Regt ([12]) develops Feynman’s view that understanding a formalism is a matter of seeing what the solution to a problem will be without having to compute it explicitly. But as de Regt points out,

¹⁸Noncommuting time is a delicate issue, technically and conceptually. See [26, 3] for discussion.

while our geometric intuitions are a fruitful resource for ‘seeing solutions’, they need not be the only one; again, familiarity with an algebra also allows one to anticipate when algebraic relations hold without explicit calculation.

So there do not seem to be in principle barriers to developing an algebraic theory. All the same, commutative space seemingly provides a very useful tool for investigating ontology, insofar as individuals can be separately localized, and the parts of space give a way of distinguishing parts of individuals. (Of course, even non-relativistic quantum non-locality makes this road to ontology treacherous at best.) Or again, *physical* reality is often tied to causal connectedness, which in turn is most readily understood in terms of effects propagating in space. So despite our metaphysical sophistication it is still puzzling to know where to start in talking about the ontology of a theory like NCG, in which the familiar spatial handles are missing. What, then is there? How can we discern a coherent ontology from the theory?

Since we are considering a theory that replaces differential geometry with algebra it will be useful for the following to bear in mind two kinds of interpretational moves made regarding space. There are two main questions at stake: (i) to what extent, if any, are the points of a differential manifold real, physical objects, akin to material systems? (ii) What aspects of spatiotemporal structure, such as topological and geometrical relations, are fundamental (capable, for instance, of providing ‘deep’ explanations)? Especially, following the (re-)introduction of the ‘hole argument’ ([18]), the locus of philosophical debate was on the first question: the ‘manifold substantialist’, who holds the points to be physically real, is faced with versions of Leibniz’s shift arguments, in which one imagines the material content of the universe rearranged in spacetime. Earman and Norton’s argument makes the point especially sharp in theories with dynamical geometries, such as GR, since then the problem of indeterminism can (arguably) be added to that of underdetermination: because of diffeomorphism symmetry there are solutions of the Einstein field equation that agree up to a given Cauchy surface, but differ by a diffeomorphism after.

Recent work has largely focussed on the second question. The idea of ‘dynamical interpretations’ of spacetime theories is that certain spatiotemporal structures (particularly affine and metrical ones) are not fundamental, but merely represent, say, the symmetries of the laws of material systems; hence ‘real’ explanations (of time dilation, for instance) are in terms of how systems behave according to physical laws, not geometry (for instance, ([5]); such interpretations are discussed in ([25, 6])). With all this in mind, consider some possible interpretational strategies; we do not aim to adjudicate between them, but merely demonstrate the possibility of an algebraic interpretation.

First, in the formalism of NCG, instead of spatial points and their relations, we have elements of an algebra and their relations; this observation suggests that the elements could be thought of, metaphysically, along the lines of points. To pursue this idea more concretely let’s take the algebra \mathcal{A} to be \mathcal{R}_θ^d , the algebra of polynomials of the noncommuting base elements, \hat{x} and

\hat{y} .¹⁹ ‘Algebraic substantivalism’ then attributes to the elements of \mathcal{R}_θ^d the same kind of ‘physical reality’ that manifold substantivalism attributes to points. To be a little more careful, just as the latter view takes mathematical points to represent, more-or-less literally, physical points, so algebraic substantivalism takes the elements of the mathematical algebra to represent, more-or-less literally, physical objects, which we shall continue to call ‘fields’. To be clear, the mathematical representation of the NCG is not itself something physical, but, according to substantivalism, what it represents is.

The idea that the points of a mathematical spacetime manifold could represent points of physical spacetime seems to be a natural one; at least philosophers (including Newton and Leibniz) have taken it (or something like it) to be a view worth defending or disputing. It appears that applying parallel reasoning in the parallel case of NCG is feels less natural. However, the only differences between the two cases lies in the non-spatiality of the fields. But that is no reason to reify in one case and not the other: as far as existence goes, we have already rejected spatiality as a condition. And while non-spatiality makes the fields less immediately connected to objects of experience, we shall see below how NCG does connect with experience. In other words, however manifold substantivalism views points, algebraic substantivalism views elements of the algebra; understand one and you understand the other.

John Earman ([17], §9.9) proposed using such an interpretation to advance the substantivalist-antisubstantivalist debate, based an algebraic formulation of general relativity, known as an ‘Einstein algebra’, due to Robert Geroch [24]. Because an Einstein algebra fixes a spacetime only up to diffeomorphism, it seems (but see below) that symmetric situations get the very same algebraic descriptions; so Earman suggested that an interpretation that takes the algebra as fundamental would thus avoid the hole argument. Moreover, Earman described the interpretation as substantival ‘at a deeper level’.

For an alternative interpretation, suppose one has a noncommutative field theory (NCFT) of a scalar field: suppose that the dynamical object of the theory is a scalar field, as in electromagnetism one has a theory of an antisymmetric tensor field. For want of a better term, call this a ‘material field’, to distinguish it from the fields of the algebraic geometry (though it is only ‘material’ in the same sense that the electromagnetic field is). Then, each (algebraic) field corresponds to a *possible* state of the material field, and can be interpreted to *be* such. This move is reminiscent of the anti-substantivalist proposal that points are merely possible locations, not physical objects themselves; the typical response to this suggestion is that it simply introduces new entities with all the troubling features of points, and so the difference is too small to generate a truly distinct view in the interpretation of spacetime theories. But note that in scalar NCFT things are potentially more promising in that no new possibilities are proposed, because the material field is already assumed to be a determinable, with many possible determinate ‘configurations’, with a noncommutative structure. Of course one might say that locations are also possible states of spacetime objects, but

¹⁹Standing in for physically realistic algebras of more complex objects like tensors and spinors.

in that case there is the option of taking *relative positions* to capture locations. There seems to be no corresponding move for the states of a noncommutative scalar field.²⁰

A third option for interpretation is suggested by Bain ([1]) in the context of the Einstein algebra formulation of general relativity. In that case diffeomorphisms correspond to automorphisms of the algebra, and so Leibniz shifts have an algebraic counterpart: there are maps of the algebra onto itself that preserve the algebra, and hence the geometry. Do we not [34] recapitulate hole indeterminacy at the algebraic level? Not, it seems, if pose the question of determinism as follows: for any two algebraic models whose representations, restricted to $t < T$, are the same, are they the same representations *tout court*? Given the representation theorems about Einstein algebras, the answer is ‘yes’. However, Bain, is unconvinced by this response. He proposes that the fields only have identities in virtue of their algebraic relations to one another – a version of ‘structuralism’, since the fields become bare *relata* for the essential, algebraic, content of the theory. In this case, since the structure is preserved by automorphisms, such shifts make no difference; the analogy to similar moves in the hole argument – or in response to other issues arising from shifts – should be clear. One even (in principle) could go a step further, and treat the elements of the algebra as purely formal, not representing any physical ‘fields’, even with weakened identity conditions. What would be left would be pure structure. It seems that all of these structuralist moves could be made just as well in the noncommutative as commutative cases (though the question of indeterminism remains).

Algebraic substantivalism and structuralism, and the idea that fields are states of a scalar field focus on the ontology of the elements of the algebra, as similar views concern the status of points in standard spacetime theories, but perhaps one could also attempt a ‘dynamical’ interpretation of NCG. Such a view would take the ‘material’ noncommuting field as the fundamental thing, and view the algebra (certainly the algebraic relations, but potentially also its elements) as merely representing something about its equations of motion. How such a view might be developed, how plausible it might be, and how it might relate to the others described here, are questions that will have to be addressed at another time.

5 Recovering space from NCG

All of the mathematical generalisations, and suggestive analogies might be of only abstract interest, if it weren’t the case that the framework of modern *physics*, also survives the transition to noncommutativity. The geometry and calculus involved can be given algebraic form, and survive. Importantly, the Lagrangian that characterizes any theory, can also be fully rendered in algebraic terms, even in the noncommutative case: so we have physics in a noncommutative geometry, specifically, a ‘noncommutative field theory’ (though note, as in §3, it is the geometry that the fields

²⁰([15]) proposes an approach along these lines for spacetime in ordinary QFT, in which points are identified with sets of local observables, themselves taken as possible properties of systems.

‘inhabit’ that is noncommutative, as the fields themselves have not been quantized at this stage). Moreover, other important pillars of modern physics, like Noether’s Theorem, also survive (it only requires that the algebra be associative): hence the central importance of conserved currents remains. (And gauge fields also exist, though importantly the distinction between ‘internal’ gauge symmetries, and ‘external’ spatial symmetries is blurred.) In this section, we discuss a proposal for the recovery of the manifest image of a classical (commuting) space from a noncommutative field theory. We begin, in §5.1, by introducing two formally equivalent representations of a noncommutative field theory. Then, in §5.2 we discuss and assess a proposal about the emergence of ordinary spacetime from a this algebraically construed, fundamentally noncommutative field theory. We conclude in §5.3 with a short reflection on the problem of empirical incoherence that we touched upon earlier.

5.1 The Moyal and Weyl representations

Let us try to understand how spacetime might emerge from a theory of non-spatial degrees of freedom. On the one hand, *prima facie* we have no degrees of freedom that are intuitively ‘spatial’: no point-valued fields, but instead the state-values are simply elements of the algebra \mathcal{R}_θ^d . While on the other, the theory apparently contains enough structure that one successfully might connect it (in some kind of limit) with familiar physics, in a classical, phenomenal, spacetime. (Note that in this section, the fields, while noncommuting, are still not fully quantized: we do not have \hbar -featuring commutation relations on the canonical variables. We turn to the fully quantum version in §5.2.)

Earlier (§3.2), we gave an operator representation of the noncommuting algebra, but we can equivalently represent it in terms of functions on a manifold, by deforming pointwise multiplication to a new product. For instance, in our toy case, with the algebra \mathcal{R}_θ^d , we take the usual smooth coordinate functions x, y , etc as base elements, and introduce the Grönewold-Moyal—or simply Moyal—product:²¹

$$\phi(x) \star \psi(x) = \phi(x) \cdot \psi(x) + \sum_{n=1}^{\infty} \left(\frac{i}{2}\right)^n \frac{1}{n!} \theta^{i_1 j_1} \dots \theta^{i_n j_n} \partial_{i_1} \dots \partial_{i_n} \phi(x) \cdot \partial_{j_1} \dots \partial_{j_n} \psi(x). \quad (8)$$

Clearly the \star -product contains new terms in addition to ordinary multiplication.²² One just needs to observe that the new terms form an (infinite) sum of derivatives with respect to the coordinates,

²¹See [35] for a full treatment. Here we continue to assume flat, infinite space, but the representation extends to more complex cases. We also restrict attention to fields that vanish smoothly at infinity (so that the ‘physicists fundamental theorem of the calculus’ reads $\int dx df(x)/dx = 0$). This restriction is a common but notable assumption in physics: on the one hand it is justified locally by the assumption that arbitrarily distant differences are irrelevant; on the other hand it raises questions about the universality of physical theories.

²²Note that this star operation has nothing to do with the involution operation on \star -algebras. The product is not unique, and there are several others which reproduce the commutation relation (6) (all of the translation invariant one are described in [22]).

5.1 The Moyal and Weyl representations

weighted by the elements of θ . For instance,

$$\hat{x}\hat{y} \rightarrow x \star y = xy + \frac{i}{2}\theta^{xy} \neq xy. \quad (9)$$

We can use (9) to verify the relevant commutator in the Weyl transform:

$$[x, y]_{\star} \equiv x \star y - y \star x = xy - yx + \left(\frac{i}{2}\right)\theta^{xy} - \left(\frac{i}{2}\right)\theta^{yx} = i\theta^{xy}, \quad (10)$$

by the antisymmetry of θ^{xy} . Thus commuting coordinates do indeed have the algebra of the noncommutative theory, *with respect to the Moyal product*.

Another instructive example is the product of Gaussians, which we might take to represent (as before) a localized ‘particle’ ρ and measuring device g . Supposing both have width α , and locations x_1 and x_2 , respectively, the product is:

$$g \star \rho = \frac{\alpha^4 \exp\left(-\frac{\alpha^2(x_1^2+x_2^2)-2x_1(\alpha^2x+2i\theta y)-2x_2(\alpha^2x-2i\theta y)+2\alpha^2(x^2+y^2)}{\alpha^4+4\theta^2}\right)}{(\alpha^4 + 4\theta^2)} \quad (11)$$

Note that the product is not commutative because of the presence of the phase $2i\theta y(x_1 - x_2)$ which is not symmetric under the exchange $x_1 \leftrightarrow x_2$, therefore $g \star \rho \neq \rho \star g$. Moreover, the result, which one would like to interpret as a probability, is a complex number. This latter aspect could in principle be resolved taking the modulus, but it is clear that the ‘classical’ interpretation is becoming untenable.

We can also compare it with the product of classical Gaussians, to see the difference that non-commutativity makes. Let the Gaussians be sharply peaked, with $\alpha = 0.1\sqrt{\theta}$, at the distance $0.5\sqrt{\theta}$. In the classical case the ordinary commutative product is practically zero everywhere (too small to be plotted), and the expectation value $\langle x^2 + y^2 \rangle$ is effectively zero. However, things are quite different for noncommuting fields, and the star product is nonvanishing, and quite spread, as can be seen in figure 1. Although the integral which would give rise to the expectation value in this case does not change,²³ all of the other momenta of the probability change, for example

$$\langle (x^2 + y^2)g \rangle_{\rho} = \frac{e^{-\frac{(x_1-x_2)^2}{2\alpha^2}} (\pi\alpha^4 (2\alpha^2 + (x_1 + x_2)^2) - 4\pi\theta^2 ((x_1 - x_2)^2 - 2\alpha^2))}{8\alpha^2}$$

which is not positive definite.

In other words we see again the impossibility of operationalizing position measurements below the noncommutativity scale: even though the ‘particle’ is supposedly five times the resolution of the device away, it overlaps with the device, though *no* probability can be assigned to its detection. From the manifold point of view there is no appreciable pointwise overlap of the functions, so this effect appears to be non-local. Better, we should (again) recognize that we have a failure to

²³This is a particularity of the Moyal product as opposed to other noncommutative geometries.

5.1 The Moyal and Weyl representations




Figure 1: *The real and imaginary parts of (11) for width $0.1\sqrt{\theta}$, at the distance $0.5\sqrt{\theta}$.*

operationalize position measurements below the noncommutativity scale, and that hence there really are no smaller regions, and no points, once the geometry is noncommutative.

Working in the Moyal star (or ‘Weyl’) representation greatly facilitates extracting the physical consequences of the theory because the usual methods of the calculus (and hence of standard field theory, including QFT) can then be applied. One ‘simply’ has to multiply physical quantities, not in the usual way, but with the \star -product: an area is $x \star y$ (or the unequal $y \star x$) not $x \cdot y$; fields given as series expansions of the x^i s are to be understood in terms of expansions using the \star -product (for instance, exponentials); and terms in the equations of motion involve \star -multiplication. In other words, for every physical purpose, the \star -product is the relevant operation, and ordinary multiplication is only relevant insofar as it is involved in the definition (8) of the putatively ‘real’ multiplication. In yet other words, the physical facts don’t care about the commutativity of space, and it is thus natural to see it just as a convenient way of representing, the real noncommutative nature of space.

So it would then be a mistake to think, because there is such a representation, in terms of smooth functions on a manifold, that after all noncommutative geometry is a theory of perfectly ordinary space. As we have been at pains to show, the notions of arbitrary small regions and of points are not well-defined in any sense in such a theory. In the *Weyl representation* of the algebra, the manifold is exactly that, a component of the representation, with excess representational structure for the true, essentially algebraic, fundamental objects.

It’s important to stress at this point that although elements of the algebra are represented by *functions*, $\phi(x)$ over space, neither the value at any point, nor the restriction of $\phi(x)$ to any region corresponds to anything in the algebraic formulation, and so has any fundamental physical significance. (Let that sink in.) Normally we think of the value of a field at a given point as conveying some physical meaning, such as the electric field strength, but in the Weyl representation, this is not the case: only the full function corresponds to an element of the algebra and corresponds to something fundamental. Obviously this situation presents a puzzle, for most familiar physical

quantities are associated with points and finite regions. We turn to this puzzle next.

5.2 Finding Spacetime

The Weyl representation provides a way of extracting empirical consequences from NCG. First, one can take the equations of motion for a classical field theory in ordinary commuting space, and rewrite them replacing all ordinary multiplication with Moyal star multiplication: the result is the equations of motion for a NCFT expressed in its Weyl representation. One obtains a more complex, but otherwise formally standard field theory – though of course noncommutation of the coordinates undermines a straight-forward interpretation. At this point, we shift to an unequivocally quantum theory by formulating a ‘second quantized’ NCFT; in particular, this the machinery of the path integral formalism to be brought to bear. Finally, those standard methods allow the derivation of empirical results, especially probabilities for particles to scatter off one another in different ways – scattering ‘cross-sections’.

These are often tacitly taken by physicists (at least those working in the QFT program) as basic concrete—indeed empirical—spacetime objects. That is, scattering occurs at an (extended) location so gives meaning to place; and depends, amongst other things, on the metric at that location, so gives meaning to geometry. Philosophically speaking, cross-sections are, crudely speaking, material objects to which a relationist might attempt to reduce space; or perhaps better, from which one might give a ‘dynamical interpretation’ of spacetime. We don’t take any stance on whether physicists have such programs in mind, but rather draw on the common tacit assumption in, for instance, string theory that our experiences of space can be recovered through scattering cross-sections, from a theory that is fundamentally not spatial. Again, the idea being that scattering gives empirical content to extended location and geometry; so recover a set of scattering amplitudes, and you have recovered spacetime. This construction is discussed in more detail in ([28, chapter 7]).

In particular, consider a world in which the cross-sections of a NCFT turn out to be correct. To understand the place of observed, classical space in such a NCFT it therefore makes good sense to focus on these cross-sections; there are other empirical aspects, and other ways in which space can be found, but cross-sections exemplify both perfectly. So we will proceed (and indeed speak) as if to understand the meaning of cross-sections in NCFT *is* to understand classical spacetime. More could be said, but we expect it to be more of the same, and not to add to the central issue. And of course, we need to a way to understand ‘pointy’ classical spacetime because we have argued that points and small regions have no physical meaning; and especially if one adopts an ontology in which the algebra is fundamental, for as we noted then neither points nor regions have fundamental significance. Our proposal here, then, is that recovering empirically meaningful cross-sections is to give derivative, empirical meaning to ordinary spacetime: to show how it ‘emerges’. (Though without reifying points.) This question has been addressed by Chaichian, Demichev and Presnajder (hereafter CDP) in a very interesting paper ([8], what follows is based

on their analysis, though it suggests a somewhat different solution.

The problem of finding scattering cross-sections can be further reduced to the calculation of ‘2-point functions’: squared, these represent the probability that, left to itself, a quantum at x in space and time would be ‘found’ at y , the simplest kind of ‘scattering’.²⁴ These, along with interaction terms, are the ingredients of the Feynman method for calculating cross-sections, so they can be taken as giving the empirical spatial content of a QFT—and hence of NCFT. The 2-point functions make the problem of giving a spatial interpretation very clear, for they are functions of x and y , coordinates in phenomenal, commuting spacetime – and so have no immediate significance in NCFT, in which the coordinates cannot be ordinary number-valued, since they don’t commute! That is, finally, the question of the meaning of phenomenal cross-sections—so of *space*—in NCFT narrows to the question of the significance of the *commuting* coordinate arguments of the 2-point functions.

Let us take a closer look at the 2-point functions: these are the vacuum state expectation values of a product of field operators—equation (8). As we have emphasized, to this point we have not quantized the fields (in the narrow sense) by imposing a noncommutative algebra. Now we do take that step: take the state of the unquantized field, $\phi_W(x)$, which takes as its values field configurations in the Weyl representation; and second quantize it, promoting it to a quantum operator $\hat{\phi}_W(x)$. The corresponding 2-point function, $G_W(x_1, x_2)$, can be written

$$G_W(x_1, x_2) = \langle 0 | \hat{\phi}_W(x_1) \hat{\phi}_W(x_2) | 0 \rangle. \quad (12)$$

We quantize using the path integral formalism, in which such quantities are given by field integrals over the classical fields, weighted by the action:

$$= \int D\phi_W \phi_W(x_1) \phi_W(x_2) \cdot e^{i\hbar \int d^n x \mathcal{L}(\phi_W(x), \dot{\phi}_W(x))}. \quad (13)$$

Note that because we have fully quantized, \hbar has finally made an appearance.

This expression makes the interpretational issue very clear, for the dependence on the field at x_1 and x_2 is explicit in the (functional) integral. But, as discussed at the end of §5.1, the point-value of a field in the Weyl representation has no fundamental meaning – only the field configuration over the whole space represents anything in the algebraic formulation, namely an element of the algebra. So the same is true of the 2-point function: it can have some significance as a function over $\mathcal{R}^d \times \mathcal{R}^d$, but its point-values, or its integral over a region, do not. But these are exactly what we would like to take as scattering amplitudes, the empirical content of the theory.

A first response would be to more-or-less ignore this situation. One simply takes the coordinates in the Weyl representation to correspond to ‘phenomenal coordinates’—the ones by which we label points of ordinary, observed space. At first glance it looks as if this response simply throws

²⁴Saying what ‘found’ means in such contexts is to propose a solution to the measurement problem, something we are trying to sidestep as far as possible.

away the noncommutative spacetime and views the theory as one with unusual equations of motion; if the coordinates are just those of ordinary commuting space, then we just have a QFT in that space with a standard lagrangian modified by use of the Moyal star. However, while this approach might be expected to produce decent predictions over distances above greater than $\sqrt{\theta}$, it is conceptually incoherent in virtue of being undefined at distances less than $\sqrt{\theta}$, as we showed in §2–3. Once again, the ‘pointlessness’ of noncommutative geometry is the main point of this paper!

A second response, which recognizes this situation, to recover the appearance of commuting spacetime, is that based on the work of CDP [8]. To keep things simple, we work in two dimensions, with $[\hat{x}_1, \hat{x}_2] = \theta$. As we saw, because of this noncommutivity, the Weyl field operators do not have the usual interpretation as localized quantities, but that doesn’t mean that the same is true for other operators in the theory. Indeed, we should expect that some other observables do represent empirical spacetime quantities.

In particular, we will consider Weyl fields that are ‘smeared’ over a region of order θ ; the idea being that these are insensitive to sub- θ physics, while capturing the physics of super- θ regions. More specifically, and in the spirit of §3, we propose that the following maps Weyl fields into the commuting fields of familiar empirical spacetime physics: i.e., fields describing observed scattering phenomena.²⁵

$$\phi_P(x) \equiv \int d^2x' \frac{e^{-(x-x')^2/\theta}}{\pi\theta} \phi_W(x'). \quad (14)$$

Smearing fields is a common practice in QFT, usually done to avoid pathological behaviours at short distances. Here we elevate it to a way to operationally use an object, a point, as an approximate avatar to connect with classicality.

This proposal is a ‘guess’, a hypothetical part of the theory, subject to testing, and potentially to replacement by some other ansatz; but it is based on the most natural way of relating noncommuting and commuting space. (Note too that the form of the smearing is the simplest, rotationally invariant form one can have.) But we can take the x -arguments of these ‘new’ fields to be those of observed space; while x' is the coordinate of the space in which the Weyl transforms live. That is, (14) can be read as a map into the reals, from observed, commuting space and Weyl field configurations (the integral means that the map depends on the full configuration): $x \times \phi_W(x') \rightarrow \mathcal{R}$.

Given (14) and the interpretation of $\phi_P(x)$, the 2-point function for the phenomenal fields is given by the path integral prescription:

$$G_P(x_1, x_2) \equiv \langle 0 | \hat{\phi}_P(x_1) \hat{\phi}_P(x_2) | 0 \rangle \quad (15)$$

$$= \int D\phi_W \phi_P(x_1) \phi_P(x_2) \cdot e^{i \int d^n x \mathcal{L}(\phi_W(x), \dot{\phi}_W(x))} \quad (16)$$

²⁵Technically, these fields are the ‘normal’ symbols of the noncommuting fields.

which is simply the smeared version of the Weyl 2-point function:

$$= \int d^2x'_1 d^2x'_2 \frac{e^{-(x-x'_1)^2/\theta}}{\pi\theta} \cdot \frac{e^{-(x-x'_2)^2/\theta}}{\pi\theta} G_P(x'_1, x'_2). \quad (17)$$

Note that at this point that we diverge from CDP proposal. Their idea is that the action in the path integral should be rewritten in terms of the phenomenal field ϕ_P . Their approach amounts to treating the phenomenal field as the true degrees of freedom. Instead, what we suggest is that we treat the Weyl fields as the true degrees of freedom, as we should if we take the noncommuting spacetime seriously: we simply recognize that the canonical degrees of freedom are not those we experience as phenomenal fields – those are represented by ϕ_P . Again, that hypothesis (in conjunction with the rest of the theory) is testable, and links the fundamental theory to experiment.

A puzzle arises, for if $\phi_P(x)$ has physical—if derivative—significance, then it seems as if its point values, and values over sub- θ regions do too; contrary to everything we have argued! But of course it is part of our proposal that they do not; only the differences over super- θ regions have physical meaning. The point values of $\phi_P(x)$ can only be understood as representational baggage, required by the formalism that we have adopted to formulate NCFT, and connect it to experience. (Of course, smearing enforces this interpretation, since it assumes the irrelevance of sub- θ physics.) Recall that the puzzle we are trying to solve is how to assign physical significance to *any* regions if the fundamental objects are essentially algebraic, and non-spatiotemporal. The CDP ansatz addresses that issue, without giving significance to points.

One could complain that the *theoretical* meaning of the phenomenal field is unclear – the $\phi_P(x)$ can formally be defined according to (14), but can we get a clearer insight into their place in the theory? In particular, do the phenomenal x s have an interpretation in the theory, since they are not the noncommuting coordinates? Since they label points of phenomenal space, an answer will illuminate how phenomenal space is found in a NCFT. CDP suggest an answer ([8],): they note that the phenomenal fields are equal to the expectation values of the noncommuting fields in so-called ‘coherent’ states, $|\xi_x\rangle$ ²⁶:

$$\phi_P(x) = \frac{\langle \xi_x | \hat{\phi}_W | \xi_x \rangle}{\langle \xi_x | \xi_x \rangle} \quad (18)$$

Coherent states are the quantum states with a semi-classical behaviour. Here they take the added role of giving an operational meaning to the points of spacetime, albeit in their smeared guise. In the Weyl representation, a coherent state can be thought of as an isotropic state, centered on a point, x ; the x s can be taken as their quantum numbers. Then (18) shows how the point-values of the phenomenal field can also be understood as labelled by coherent states, taking the point in phenomenal space to be the corresponding quantum number.²⁷

²⁶Specifically, $|\xi_x\rangle = \exp(x_1 + ix_2)a^\dagger|0\rangle$

²⁷CDP approach things from the other direction. They propose that the phenomenal fields should be

But we will end the discussion here, having suggested how algebraic interpretations might be given, and shown how to recover the appearance of commutative spacetime, but with fields with no meaning below a certain scale—so again, pointless physics.

5.3 Empirical coherence and physical salience

More generally, Maudlin[32] questions the feasibility of ‘deriving’ classical spacetime from some non-spacetime theory (he has in mind deriving 3-space from $3N$ -configuration space, but the point generalizes). At the heart of his concern is that even if a formal derivation can be found, involving a mathematical correspondence between classical spacetime structures and structures defined in terms of a (more) fundamental non-spatiotemporal theory, it does not follow that the classical spacetime just *is* the more fundamental structure. Mathematical correspondences are too cheap: for instance, many systems are described by simple harmonic oscillator equations, but it would be a mistake to conclude that they were physically indistinguishable just because of this formal correspondence. According to Maudlin, for a reductive account, a formal derivation must also be ‘physically salient’. We take this to mean that the derivation must veridically track the way in which fundamental structures ‘combine’ to physically constitute derivative ones. For instance, in ideal gas theory the *formal definitions* of ‘temperature’ as mean kinetic energy and ‘pressure’ as momentum transfer *track* the corresponding phenomenal thermodynamical quantities: kinetic energy is transferred between the molecules of the gas to liquid in a thermometer causing its expansion; and the pressure on the side of a vessel is due to the molecules contained colliding with it. The problem with a fundamental theory without spacetime is that our notions of what kinds of derivation track in this way are spatiotemporal notions, relying on colocation and dynamical interaction (of gas molecules with thermometers or vessel walls, say), for instance. But such familiar notions cannot apply if the physics involved is not, by supposition, itself spatiotemporal. So we face two problems: first, what new notions might apply? And second, even if we have a proposal, on what grounds can we conclude that we are correct?

If the analysis of this paper is correct, then noncommutative geometry is a nice example of this situation: the fundamental structure is algebraic, not a commutative geometry, and so concepts like ‘spatial location’ are not primitives of the theory. Rather, spatial structure is derived. In particular, we have discussed the proposal that it be recovered via CDP ansatz, which we have also argued is not entailed by the theory, but an additional postulate. More precisely, it is an interpretational postulate, specifying how algebraic objects can ‘combine’ to physically constitute classical spatial structures – *a novel proposed non-spatiotemporal conception of which derivations are physically salient in the theory*. Thus the first problem can has been addressed in this case. (As in most cases, there is enough spatiotemporal structure in the underlying theory – which is after all a deformation of a commutative geometry – to find clues about how to reconstruct spacetime.)

As for the second problem, [27] proposes that such postulates, concerning how more funda-
labelled by the coherent states, and then conclude that they should be smeared according to (14).

mental structures compose to constitute less fundamental ones, are justified a posteriori, not a priori. (The paper briefly discusses NCFT and with other examples of theories without spacetime, along the lines found in this paper: identifying what spatiotemporal features are lost in each case, and explaining how they may be derived.) That is, how the fundamental gives rise to the less fundamental is not a matter of metaphysical necessity, but of physical contingency, and so is something that can only be discovered empirically, with the theory itself. For instance, if a theory of noncommutative geometry was empirically successful (in the usual ways, especially in making novel predictions that cannot be accounted for in any other known way) then both the theory *and* CDP interpretational ansatz would be confirmed. That is, ultimately we justified in believing a derivation to be physically salient in the same way that any other scientific belief is justified: through successful confrontation with the data. No more is possible, but then it never is.

Thus, in addition to introducing NCFT and raising some specific interpretational questions, this paper presents it specifically as an example of derived or ‘emergent’ space, in order to illustrate and address Maudlin’s challenge. There is a gap between noncommutative and commutative geometries, which can be *formally* filled by the CDP ansatz; but if this strategy were empirically successful then we would have scientific grounds to further believe that the derivation is physically salient, that the ansatz is a veridical statement of physical composition. The hope is that working through the example makes that claim plausible, or at least intelligible.²⁸

6 Conclusion

This paper had three goals beyond introducing noncommutative geometry to an audience of philosophers. The first was to convince the reader that one of the physical upshots of a theory having a noncommutative geometric structure is that it cannot include, in its ontology, a set of arbitrarily small spatial regions (*a fortiori* points). We demonstrated this in two ways: in terms of the undefinability of arbitrarily small distances, using Connes’ spectral triples, and in terms of the non-operationalisability of arbitrarily small separations, using measuring devices modelled by Gaussians in space. The second was to propose an appropriate ontology for a field theory set in a noncommutative space. We suggested that a field-first ontology was the only sensible option, and explored some of the consequences of this proposal. We were then led by this commitment to a puzzle about how to account for the appearance of a commutative spatiotemporal geometric structure, at least at the level of experience and experimental data. The final goal, then, was to discuss and assess a dynamical proposal, originally due to Chaichian, Demichev and Presnajder [8] for the emergence of spacetime using the resources of our best theory of matter–quantum field theory.

²⁸Of course it is logically possible to deny it, but we would say (a) that the historical record contains examples of similar changes in the concept of physical salience, and (b) that such a denial would collapse into a general antirealism, which is not our target here – rather the issue is whether there are special reasons to dispute the physical salience of derivations of spacetime.

As promised in the introduction, we demonstrated, via considerations of noncommutative geometries, one way of embedding a class of extant, well-confirmed physical theories (in this case, quantum field theories) in a broader logical landscape. This is, of course, a small step in the direction of unearthing all the important tacit commitments associated with interpretations of such theories. Even so, we have made some real progress on that front. We realised that we can, in fact, talk about spacetime without it being indiscrete.

References

- [1] Jonathan Bain. Einstein algebras and the hole argument. *Philosophy of Science*, 70(5):1073–1085, 2003.
- [2] Jeffrey A Barrett. *The quantum mechanics of minds and worlds*. Oxford University Press, Oxford, 1999.
- [3] H. Bozkaya, P. Fischer, H. Grosse, M. Pitschmann, V. Putz, M. Schweda, and R. Wulkenhaar. Space-time noncommutative field theories and causality. *Eur. Phys. J. C*, 29:133–141, 2003.
- [4] Percy Williams Bridgman. *The logic of modern physics*, volume 3. Macmillan New York, 1927.
- [5] Harvey R Brown. *Physical Relativity*. Oxford University Press, Oxford, 2005.
- [6] Harvey R Brown and James Read. The dynamical approach to spacetime. In Eleanor Knox and Alastair Wilson, editors, *The Routledge Companion to Philosophy of Physics*. Routledge, forthcoming.
- [7] Claudio Calosi and Jessica Wilson. Quantum metaphysical indeterminacy. *Philosophical Studies*, 176(10):2599–2627, 2019.
- [8] Masud Chaichian, A Demichev, and P Prešnajder. Quantum field theory on non-commutative space-times and the persistence of ultraviolet divergences. *Nuclear Physics B*, 567(1-2):360–390, 2000.
- [9] Alain Connes. *Noncommutative geometry*. Academic Press, Cambridge, MA, 1995.
- [10] George Darby. Quantum mechanics and metaphysical indeterminacy. *Australasian Journal of Philosophy*, 88(2):227–245, 2010.
- [11] George Darby and Martin Pickup. Modelling deep indeterminacy. *Synthese*, forthcoming.
- [12] Henk W De Regt. Spacetime visualisation and the intelligibility of physical theories. *Studies In History and Philosophy of Science Part B: Studies In History and Philosophy of Modern Physics*, 32(2):243–265, 2001.

References

- [13] Jan Dereziński and Maciej Karczmarczyk. Quantization of gaussians. *arXiv preprint arXiv:1701.07297*, 2017.
- [14] Neil Dewar. Sophistication about symmetries. *The British Journal for the Philosophy of Science*, 70(2):485–521, 2019.
- [15] Dennis Dieks. Space and time in particle and field physics. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics*, 32(2):217–241, 2001.
- [16] John Earman. Leibnizian space-times and leibnizian algebras. In *Historical and Philosophical Dimensions of Logic, Methodology and Philosophy of Science*, pages 93–112. Springer, 1977.
- [17] John Earman. *World Enough and Space-Time: Absolute versus Relational Theories of Space and Time*. MIT Press, Cambridge, MA, 1989.
- [18] John Earman and John Norton. What price spacetime substantivalism? the hole story. *British Journal for the Philosophy of Science*, pages 515–525, 1987.
- [19] Herbert Enderton. *A mathematical introduction to logic*. Elsevier, 2001.
- [20] Benjamin H Feintzeig. On theory construction in physics: Continuity from classical to quantum. *Erkenntnis*, 82(6):1195–1210, 2017.
- [21] Hartry Field. Can we dispense with space-time? In *PSA: Proceedings of the Biennial Meeting of the Philosophy of Science Association*, volume 1984, pages 33–90. Philosophy of Science Association, 1984.
- [22] Salvatore Galluccio, Fedele Lizzi, and Patrizia Vitale. Twisted Noncommutative Field Theory with the Wick-Voros and Moyal Products. *Phys. Rev. D*, 78:085007, 2008.
- [23] I.M Gelfand and M.A Naimark. On the embedding of normed rings into the ring of operators in hilbert space. *Mat. Sbornik*, 12:197–213, 1943.
- [24] Robert Geroch. Einstein algebras. *Communications in Mathematical Physics*, 26(4):271–275, 1972.
- [25] Nick Huggett. Essay review: Physical relativity and understanding space-time. *Philosophy of Science*, 76(3):404–422, 2009.
- [26] Nick Huggett, Tiziana Vistarini, and Christian Wüthrich. Time in quantum gravity. In Adrian Bardon and Heather Dyke, editors, *The Blackwell Companion to the Philosophy of Time*, pages 242–261. Blackwell, Chichester, 2013.
- [27] Nick Huggett and Christian Wüthrich. Emergent spacetime and empirical (in) coherence. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics*, 44(3):276–285, 2013.

References

- [28] Nick Huggett and Christian Wüthrich. *Out Of Nowhere*. Oxford University Press, forthcoming.
- [29] Klaas Landsman. *Foundations of quantum theory: from classical concepts to operator algebras*. Cham: Springer International Publishing, 2017.
- [30] Lorenzo Leone. Quantizzazione di stati gaussiani. Tesi di Laurea in Fisica, 2017.
- [31] Fedele Lizzi. Noncommutative spaces. In *Noncommutative Spacetimes*, pages 89–109. Springer, 2009.
- [32] Tim Maudlin. *The metaphysics within physics*. Oxford University Press, Oxford, 2007.
- [33] Eduard Prugovecki. *Quantum mechanics in Hilbert space*, volume 92. Academic Press, 1982.
- [34] Robert Rynasiewicz. Rings, holes and substantivalism: On the program of Leibniz algebras. *Philosophy of Science*, pages 572–589, 1992.
- [35] Richard J Szabo. Quantum field theory on noncommutative spaces. *Physics Reports*, 378(4):207–299, 2003.
- [36] Jessica M Wilson. A determinable-based account of metaphysical indeterminacy. *Inquiry*, 56(4):359–385, 2013.