

# Transforms for the early Kerr metric

Stephen Athel Abbott

Independent Researcher, UK  
Email: gawxmax@gmail.com

---

## Abstract

The concept and usage of the word 'metric' within General Relativity is briefly described. The early work of Roy Kerr led to his original 1963 algebraic, rotating metric. This discovery and his subsequent recollection in 2008 are summarised as the motivation for this article. Computer algebra has confirmed that nominal transformations of this early metric can generate further natural algebraic metrics. The algebra is not abstract, nor is it advanced, however, these metrics have been overlooked for many years. The 1916 metric due to Schwarzschild misled Kerr into seeking a similar metric for rotation. The philosophy of astrophysics for Penrose, Hawking and others would have been very different had they used the two lost metrics.

## Keywords

Quasars, Pulsars, AGNs, Gravitation, LIGO

---

## 1. Introduction

The General Theory of Relativity (GTR) was established in 1915 and it was not until 1963 [1] that a model of a rotating astronomical mass was found. Since then researchers have tried to model these masses; pulsars, dwarf stars, neutron stars, stellar black holes and galactic black holes.

The metric may be described in several different ways. All are related to a formula for the spacetime interval between two events or rather the square of that value. The interval may otherwise be known as the line element within that separation formula. The metric tensor of the GTR may be referred to as a matrix rather than as a tensor and the word 'metric' may loosely refer to the separation formula itself.

The GTR is an implicit type of theory, in that a metric is required before one can study the dynamics of motion or mass distribution. More generally, the metric may be defined using several unknown functions of the coordinates for a selected coordinate system. The field equations of the GTR then constrain the form of the unknown functions within the metric.

A collection of metrics has become established in the community, mostly derived by mathematical transformation of the Kerr metric. This article describes two metrics that have not been found in the literature but use some of the easier transforms originating in 1963.

## 2. Motivation

It was quite appropriate and generous for the man who found the early metric to record his memory of the discovery. It was his 2008 summary [2] of the derivation of the original Kerr metric that has prompted the current work. In particular, it showed the use of complex variable theory to derive exact (algebraic) solutions to a system of non-linear partial differential equations. After this technical part of the derivation, certain easier geometric transformations lead to more familiar coordinate systems. It is also when approximations may have missed two exact solutions.

Subsequent to the study of Kerr's 2008 paper, and development of models for several metrics, a search was made for anything similar to the results. The nearest content [3] was from 2015. This paper begins with the confusion engendered by Kerr's sign changes with some reference to advanced and retarded solutions. It also indicates a reluctance to consider off-diagonal metric elements.

After the seminal work with complex variable, Kerr obtained his metric in terms of the variable,  $u$ . The separation formula was not originally split into the sum of 'base' and 'mass' terms. Sign changes have occurred in the splitting of the 1963 formula and the natural transformation,  $u = r + t$ , has been lost.

### 3. Kerr's metric recalled

The line element was later given by the base metric plus the mass term in the 2008 summary.

The base metric was given by...

$$ds_0^2 = (r^2 + a^2 * \cos(\theta)^2) * (d\theta^2 + \sin(\theta)^2 * d\phi^2) - ds_u^2 \quad (1)$$

where

$$ds_u^2 = (2 * dr + du - a * \sin(\theta)^2 * d\phi) * (du + a * \sin(\theta)^2 * d\phi) \quad (2)$$

The mass term was given by...

$$ds_M^2 = 2 * m * r * (du + a * \sin(\theta)^2 * d\phi)^2 / (r^2 + a^2 * \cos(\theta)^2) \quad (3)$$

where the 'line elements' have been named and the letter 'd' notation has been used for the differentials.

### 4. Strategy for nominal transforms

This author has created a repository on GitHub for Geometric Algebra with some applications to General Relativity. The repository uses the type of notebook generated by wxMaxima. Each notebook source file is an active document, in that it has a commentary interleaved with the calculation code and results are embedded next to that code when executed. The active documents for the two algebraic metrics have been uploaded to the open source repository together with copies of the references. The wxMaxima results have also contributed to the accuracy of the formulae within this article.

The transform strategy can now be described; for two different sequences of transformations, the base term and the mass term defined by Equations (1) to (3) were transformed separately for clarity. For each sequence the two resulting matrices were summed to give the metric for that sequence. The early Kerr metric was transformed to both a spherical system and also to a 'spherical, cylindrical' system of coordinates.

### 5. The linear transform

The spherical metric below was derived using only the transformation:  $r + u = -t$ . The differential,  $[du = -dt - dr]$  has been used to substitute for  $du$  since  $u$  itself does not occur explicitly in Equation (2). The base metric then becomes...

$$ds_0^2 = +d\theta^2 * (r^2 + a^2 * \cos(\theta)^2) + d\phi^2 * (r^2 + a^2) * \sin(\theta)^2 + ds_t^2 \quad (4)$$

where

$$ds_t^2 = -2 * a * d\phi * dr * \sin(\theta)^2 - dt^2 + dr^2 \quad (5)$$

The same substitution was applied to Equation (3) for the mass term. The denominator does not contain  $u$  or  $du$  and can be isolated for further calculations.

$$den = a^2 * \cos(\theta)^2 + r^2 \quad (6)$$

In order to avoid long quotients, the transformed mass term for the spherical metric has been displayed using the denominator,  $den$  as a multiplier.

$$\begin{aligned}
ds_M^2 * den = & +2 * a^2 * d\phi^2 * m * r * \sin(\theta)^4 \\
& -4 * a * d\phi * dt * m * r * \sin(\theta)^2 \\
& -4 * a * d\phi * dr * m * r * \sin(\theta)^2 \\
& +2 * dt^2 * m * r \\
& +4 * dr * dt * m * r \\
& +2 * dr^2 * m * r
\end{aligned} \tag{7}$$

The two formulae were then summed.

$$ds^2 = ds_0^2 + ds_M^2 \tag{8}$$

The usual vector of coordinates for the spacetime spherical system is given by  $(r, \theta, \phi, t)$ . The extraction of the coefficients into the metric matrix was consistent with a matrix row,  $(dr, d\theta, d\phi, dt)$  and its transpose. The extraction included the off-diagonal coefficients for  $d\phi * dr$ ,  $d\phi * dt$  and  $dr * dt$ , placed symmetrically about the diagonal and halved in value. In this way, the matrix factorization becomes equivalent to the sixteen term sum within the tensor separation formula.

All of the non-zero elements of the metric had the same denominator,  $den$ , except the coefficient of  $d\theta^2$ . The value of that element was  $den$  itself.

The following is an inspection of the Kerr metric in spherical coordinates. Firstly, only the diagonal terms for time and space, respectively, are examined here for zero rotation ( $a = 0$ ). This was for comparison with the Schwarzschild metric...

$$g_{44} = (-a^2 * \cos(\theta)^2 - r^2 + 2 * m * r) / (a^2 * \cos(\theta)^2 + r^2) \tag{9}$$

setting  $a = 0$ ,

$$g_{44} = (2 * m) / r - 1 \tag{10}$$

and for space...

$$g_{11} = (a^2 * \cos(\theta)^2 + r^2 + 2 * m * r) / (a^2 * \cos(\theta)^2 + r^2) \tag{11}$$

again setting  $a = 0$ ,

$$g_{11} = (2 * m) / r + 1 \tag{12}$$

The  $dr^2$  element,  $1 + 2m/r$ , is algebraic (no approximation); it only occurs in the Schwarzschild metric after the binomial approximation to the singular quotient,  $1/(1 - 2m/r)$ .

Now, even for the zero rotation case, there was a non-zero off-diagonal element. This retains a time asymmetry so the metric is stationary but not static...

$$g_{14} = (2 * m) / r \tag{13}$$

The non-rotating spherical Kerr metric is not equivalent to the Schwarzschild metric, it is something else...

$$ds_A^2 = (1 + 2 * m/r) * dr^2 + (4 * m/r) * dr * dt - (1 - 2 * m/r) * dt^2 + d\Omega^2 \tag{14}$$

where, in a typical notation,

$$d\Omega^2 = r^2 * d\theta^2 + r^2 * \sin(\theta)^2 * d\phi^2 \tag{15}$$

It now seems preferable to retain this natural algebraic metric rather than continue to use a diagonal form with its consequent surface singularity.

## 6. The non-orthogonal transform

For the spherical, cylindrical metric, the same transform strategy was followed and the sequence was similar to the previous metric. The two transformations were applied in turn to Equation (1) and then to Equation (3). The first transformation was  $r + u = -t$  and the second was  $r * \cos(\theta) = z$ . After the first of these, the resulting equations are Equation (4) for the base metric and Equation (7) for the mass term. Before applying the second transformation, it was required to form the differential of that transformation...

$$\cos(\theta) * dr - r * \sin(\theta) * d\theta = dz \quad (16)$$

Again, before the substitution of the circular functions themselves, the contribution of functions from the differential were included as...

$$[d\theta = (dr * \cos(\theta) - dz)/(r * \sin(\theta))] \quad (17)$$

Then, the substitution for the circular functions was...

$$[\cos(\theta) = z/r, \sin(\theta) = +\text{sqrt}(1 - (z/r)^2)] \quad (18)$$

Since the  $\sin(\theta)$  occurred as a square, there was no sign ambiguity due to the +ve  $\text{sqrt}()$  for the sine. The cosine lies in the interval  $[+1,-1]$ , and its sign is still present in the matrix coefficient for  $dr * dz$ , in Equation (20) below, as +ve and -ve values of  $z$ . The angle,  $\theta$  is measured from the spin axis ( $z$ -axis) in the range  $[0, \pi]$ .

The base metric in Equation (4) was transformed and the denominator was isolated.

$$\text{den}_b = r^4 * (z - r) * (z + r) \quad (19)$$

For clarity, the transformed base metric of the spherical, cylindrical system has been displayed using the denominator  $\text{den}_b$  as a multiplier.

$$\begin{aligned} ds_b^2 * \text{den}_b = & -d\phi^2 * r^2 * (r^2 + a^2) * (z - r)^2 * (z + r)^2 \\ & + 2 * a * d\phi * dr * r^2 * (z - r)^2 * (z + r)^2 \\ & - dr^2 * (a^2 * z^4 + r^6) \\ & + 2 * dr * dz * r * z * (a^2 * z^2 + r^4) \\ & - dz^2 * r^2 * (a^2 * z^2 + r^4) \\ & - dt^2 * r^4 * (z - r) * (z + r) \end{aligned} \quad (20)$$

The angular coordinate difference,  $d\theta$  does not occur in Equation (7) and the circular functions within the mass term were transformed directly.

$$\text{den}_m = r * (a^2 * z^2 + r^4) \quad (21)$$

Again for clarity, the transformed mass term for the non-orthogonal metric has been displayed using the denominator,  $\text{den}_m$  as a multiplier.

$$\begin{aligned} ds_m^2 * \text{den}_m = & +2 * m * a^2 * d\phi^2 * (r^2 - z^2)^2 \\ & -4 * m * a * d\phi * dt * r^2 * (r^2 - z^2) \\ & -4 * m * a * d\phi * dr * r^2 * (r^2 - z^2) \\ & +2 * m * dt^2 * r^4 \\ & +4 * m * dr * dt * r^4 \\ & +2 * m * dr^2 * r^4 \end{aligned} \quad (22)$$

The formulae for the base metric and the mass term were then summed.

$$ds^2 = ds_b^2 + ds_m^2 \quad (23)$$

The vector of coordinates for a spacetime cylindrical system would be given by  $(\rho, \phi, z, t)$ , however, for the spherical, cylindrical system they are  $(r, \phi, z, t)$  with a spherical radius.

The extraction of the coefficients into the metric from the sum of quotients given by Equation (23) was consistent with a matrix row,  $(dr, d\phi, dz, dt)$ . This included the off-diagonal coefficients for  $d\phi * dr$ ,  $d\phi * dt$ ,  $dr * dz$  and  $dr * dt$  to form a symmetric matrix. Denominators within the final matrix were formed from Equation (19) and Equation (21).

The remainder of this section is an inspection of the Kerr metric in spherical, cylindrical coordinates. The diagonal terms for time and space, respectively, are now examined for  $a = 0$  (zero rotation).

$$g_{44} = (2 * m * r^3)/(a^2 * z^2 + r^4) - 1 \quad (24)$$

setting  $a = 0$ ,

$$g_{44} = (2 * m)/r - 1 \quad (25)$$

The  $dt^2$  element is algebraic; it has the same coordinate zero as the Schwarzschild metric at  $r = 2 * m$ , and for the diagonal spatial element...

$$g_{11} = (2 * m * r^3)/(a^2 * z^2 + r^4) + (-a^2 * z^4 - r^6)/(r^4 * (z - r) * (z + r)) \quad (26)$$

again setting  $a = 0$ ,

$$g_{11} = (2 * m)/r - r^2/(z^2 - r^2) \quad (27)$$

Having applied the zero rotation condition, the spatial value for the Schwarzschild metric was found only in the plane  $z = 0$ . There is, in general, no coordinate singularity due to this element at  $r = 2 * m$ .

However, there is a singularity at  $r = +z$  and  $r = -z$ , all along the spin axis. The axial singularity is also present for non-zero rotations and for some other elements of the metric matrix. It is possible that active black holes may be modeled with this metric. We do not yet know whether the non-orthogonal system is creating a coordinate singularity or representing an actual singularity. It is tempting to conclude the former since for zero rotation the axis is arbitrary, however, this is a very special limiting condition.

## 7. Conclusion

Some of the history of the simulation of compact astronomical objects has been indicated. General Relativity and its plethora of spacetime metrics is the basis for such simulations. Within the era of LIGO success it seems timely to report the existence of two natural alternative metrics, when they have been found accidentally. This was especially so with reference to several physical realities...the extremely well focused axial features of some black holes (without the rather unlikely ring singularity of modern Kerr metrics) and the arbitrary nature of an axial feature within pulsars. Perhaps some future models may utilize the above metrics to predict the LIGO chirps from gravitational waves generated during the merger of various types of compact masses.

## References

- [1] Kerr, R.P (1963) Gravitational field of a spinning mass as an example of algebraically special metrics, Phys. Rev. Lett., **11**. <https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.11.237>
- [2] Kerr, R.P (2008) Discovering the Kerr and Kerr-Schild metrics, [https://arxiv.org/pdf/0706.1109\[gr-qc\]](https://arxiv.org/pdf/0706.1109[gr-qc])
- [3] Teukolsky, S.A (2015) The Kerr Metric, [https://arxiv.org/pdf/1410.2130\[gr-qc\]](https://arxiv.org/pdf/1410.2130[gr-qc])