

Ecological and Lyapunov Stability¹

James Justus
University of Texas, Austin
justus.phil@mail.utexas.edu

Draft: 10/15/06

Abstract: Ecologists have proposed several incompatible definitions of ecological stability. Emulating physicists, mathematical ecologists commonly define it as Lyapunov stability. This formalizes the problematic concept by integrating it into a well-developed mathematical theory. The formalization also seems to capture the intuition that ecological stability depends on how ecological systems respond to perturbation. Despite these advantages, this definition is flawed. Although Lyapunov stability adequately characterizes perturbation responses of systems typically studied in physics, it does not for ecological systems. This failure reveals a limitation of its underlying mathematical theory, and an important difference between dynamic systems modeling in physics and biology.

1. Introduction. Like many scientific concepts, fully adequate definitions of some ecological concepts have not yet been formulated. Ecological stability is one such concept. Proposed definitions of it are not fully satisfactory and, worse, seem incompatible (Shrader-Frechette and McCoy 1993). Although it is incorrect to conclude from this that the concept itself is problematic (Odenbaugh 2001), the multitude of incompatible definitions initiates and exacerbates conceptual confusion.

Among mathematical ecologists, ecological stability is commonly defined as Lyapunov stability, named after the Russian mathematician who first precisely defined the concept to describe the apparently stable equilibrium behavior of the solar system (Lyapunov 1892). His definition has found widespread application outside this context and is frequently used to analyze mathematical models of biological communities (Logofet 1993). May (1974) used this definition, for instance, in his influential analysis of relationships between the stability and complexity of such models.

¹ Presented at the 2006 Biennial Meeting of the Philosophy of Science Association, Vancouver, Canada. A Josephine De Kármán fellowship supported this research. Thanks to Mark Colyvan, Alexander Moffett, Eric Pianka, Mark Sainsbury, Carl Salk, and Sahotra Sarkar for helpful comments.

The definition has some clear advantages. Unlike other definitions, it integrates ecological stability into a thoroughly studied mathematical theory that has proved fruitful in many sciences, especially physics. It also seems to formalize the intuition that ecological stability depends on community response to perturbation. Despite these apparent advantages, ecological stability should not be defined as Lyapunov stability.

Sections 2 and 3 describe the concept of Lyapunov stability, its underlying mathematical theory, and show why this theory is so successful within physics. Section 4 considers the apparent advantages of the definition and illustrates how Lyapunov stability applies to mathematical models of biological communities. Section 5 argues this definition is problematic, focusing specifically on biological interpretation of Lyapunov stability. Based on this analysis, Section 6 draws some general conclusions about scientific definition and highlights an important difference between dynamic systems modeling in physics and in ecology.

2. Lyapunov Stability. To ensure sufficient generality, represent a system by a position vector $\mathbf{x}(t)$ (t represents time) in an abstract n -dimensional state space E . Assume E is governed by a vector function \mathbf{F} representing the magnitude and change of direction it induces on $\mathbf{x}(t)$. \mathbf{F} represents, therefore, the dynamics of the system represented by $\mathbf{x}(t)$. Points in E for which $\mathbf{F}=0$ are called equilibrium points, and unperturbed position vectors at such points remain stationary.

Lyapunov stability is a property of system behavior in neighborhoods of equilibria. Specifically, an equilibrium \mathbf{x}^* is *Lyapunov stable* in E_x ($\mathbf{x}^* \in E_x \subseteq E$) iff:

$$(1) \quad (\forall \varepsilon > 0)(\exists \delta > 0)(|\mathbf{x}(t_0) - \mathbf{x}^*| < \delta \Rightarrow (\forall t \geq t_0)(|\mathbf{x}(t) - \mathbf{x}^*| < \varepsilon));$$

where ε and δ are real values, $\mathbf{x}(t_0) \in E_x$ represents the system at some initial time t_0 , and ‘ $|\cdot|$ ’ designates a Euclidean distance metric on E . Informally, (1) says \mathbf{x}^* is Lyapunov stable if systems beginning in a \mathbf{x}^* neighborhood remain near it after perturbations that do not displace them from E_x . \mathbf{x}^* is *asymptotically Lyapunov stable* in E_x iff (1) and $\lim_{t \rightarrow \infty} \mathbf{x}(t) = \mathbf{x}^*$. The subspace E_x of E within which systems are (asymptotically)

Lyapunov stable is called the (*attraction*) *stability domain* of \mathbf{x}^* .

[Figure 1]

If the (attraction) stability domain is all of E , \mathbf{x}^* is (asymptotically and) globally Lyapunov stable.

In general, Lyapunov stability cannot be assessed with (1) because explicit solutions for $\mathbf{x}(t)$ can seldom be found. For this reason, scientific models often characterize $\mathbf{x}(t)$ in terms of differential equations:

$$(2) \quad \frac{d\mathbf{x}(t)}{dt} = \mathbf{F}(\mathbf{x}(t), \Omega);$$

where Ω is a set of parameters designating factors that influence system dynamics but are uninfluenced by them, and \mathbf{F} is a $n \times n$ matrix $[a_{ij}]$.² For the complicated systems modeled by (2) that scientists study, explicit solutions for $\mathbf{x}(t)$ are rarely available. Without such solutions, however, system behavior as required by (1) cannot be directly evaluated.

3. The Direct and Indirect Methods. Lyapunov (1892) recognized this difficulty and developed two methods for assessing (1) indirectly. The first ‘indirect method’ involves

² Since \mathbf{F} is not a function of t , (2) represents *autonomous* dynamic systems that do not explicitly depend on time. These models are common in science and the focus of the following analysis.

linearizing \mathbf{F} at \mathbf{x}^* . Specifically, let $\mathbf{A} = \left. \frac{\partial \mathbf{F}(\mathbf{x}(t), \Omega)}{\partial \mathbf{x}} \right|_{\mathbf{x}=\mathbf{x}^*}$ be the Jacobian matrix of \mathbf{F} evaluated at \mathbf{x}^* . The eigenvalues of \mathbf{A} determine whether \mathbf{x}^* is stable. These are scalar values λ_i such that $\det(\mathbf{A} - \lambda_i \mathbf{I}) = 0$, *i.e.* the roots of the characteristic polynomial of \mathbf{A} . Lyapunov proved \mathbf{x}^* is asymptotically Lyapunov stable iff:

$$(3) \quad \text{Re} \lambda_i(\mathbf{A}) < 0 \text{ for } i = 1, \dots, n;$$

where $\text{Re} \lambda_i(\mathbf{A})$ designates the real part of the i th eigenvalue of \mathbf{A} , λ_i . Since \mathbf{A} is only defined at \mathbf{x}^* , stability determined by the indirect method is restricted to infinitesimal neighborhoods of \mathbf{x}^* . For this reason, it is called *local stability*. The indirect method is prevalent in mathematical modeling because it is almost universally applicable: it applies to any system representable by differential equations like (2) and some difference equations (Hinrichsen and Pritchard 2005).

Stability criteria based on the indirect method such as (3) have a serious limitation: they provide no information about the extent of (attraction) stability domains. This prompted Lyapunov (1892) to develop a ‘direct method’ for evaluating (1). It involves constructing a differentiable scalar Lyapunov function $V(\mathbf{x})$ with an origin at \mathbf{x}^* (*i.e.* $\mathbf{x}^* = \mathbf{0}$) such that:

- (i) $V(\mathbf{x})$ is positive definite: (a) $V(\mathbf{0}) = 0$; (b) $V(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$; and,
- (ii) $\nabla V(\mathbf{x}) \cdot \mathbf{F}(\mathbf{x}, \Omega) \leq 0$ for all \mathbf{x} ;

where ‘ \cdot ’ designates the dot product, and ‘ ∇ ’ designates the gradient vector function. Lyapunov (1892) proved the existence of a Lyapunov function on $E_x \subseteq E$ ($\mathbf{x}^* \in E_x$) is sufficient for Lyapunov stability of \mathbf{x}^* in this region, and that with strict inequality in (ii), \mathbf{x}^* is asymptotically Lyapunov stable. This condition was also later proved necessary

(Hahn 1963). The ability to construct a Lyapunov function is thus a stability criterion. As a methodology, it is called ‘direct’ because its success depends directly upon the mathematical form of equations like (2), unlike the indirect method, which relies on their linearization, and unlike direct evaluation of (1), which requires explicit solutions to (2).

Although no general method for constructing Lyapunov functions is known, the direct method has proven to be an extremely useful tool for analyzing physical systems, especially in the classical framework governed by only Newtonian mechanics and friction. Across scientific fields, this is exceptional rather than typical. Constructing Lyapunov functions is usually extremely difficult for the predominantly nonlinear systems scientists study (Goh 1977). The reason for its utility in the classical framework is twofold. First, there are highly confirmed mathematical models describing numerous types of systems in this framework. The likelihood is therefore high that application of the direct method to these models will reveal the true stability properties of the systems they accurately represent. Second, there are certain quantities, such as total energy, that are conserved or monotonically dissipated in such systems, depending on how they are characterized (open or closed). These quantities ensure Lyapunov functions exist for models of these systems. Lyapunov, in fact, developed the Lyapunov function to generalize the classical energy concept, and his proof about its connection to stability depends essentially on energy conservation (Lyapunov 1892).

To illustrate, consider a closed particle mass system governed by a conservative force field \mathbf{G} in a frictionless Newtonian framework. The system energy, $V(\mathbf{x},\mathbf{v})$, is designated by the Hamiltonian:

$$(4) \quad V(\mathbf{x},\mathbf{v}) = \frac{1}{2}m|\mathbf{v}|^2 + U(\mathbf{x});$$

where \mathbf{x} is a position vector, \mathbf{v} is a velocity vector, m represents particle mass, $|\cdot|$ designates the magnitude of \cdot , and U is a scalar potential energy function such that $\mathbf{G}(\mathbf{x}) = -\nabla U(\mathbf{x})$. Energy conservation ensures:

$$(5) \quad V(\mathbf{x}, \mathbf{v}) = c;$$

where c is a constant real value. At an equilibrium \mathbf{x}^* , $\mathbf{v} = \mathbf{0}$, $\mathbf{G} = \mathbf{0}$ and hence $\nabla U(\mathbf{x}) = \mathbf{0}$. If, furthermore, \mathbf{x}^* is a local minimum of $U(\mathbf{x})$, an “energy difference” function $V^*(\mathbf{x}, \mathbf{v})$ can be defined such that $V^*(\mathbf{x}, \mathbf{v}) = V(\mathbf{x}, \mathbf{v}) - V(\mathbf{x}^*, \mathbf{0})$. $V^*(\mathbf{x}, \mathbf{v})$ is a Lyapunov function and \mathbf{x}^* is therefore stable. In this case, energy conservation entails (ii) from above is satisfied by equality. If, however, system energy were continually decreasing instead of conserved, depleted by friction for instance, (ii) would be satisfied by strict inequality and \mathbf{x}^* would be asymptotically stable.

Within the classical framework, stability properties of more complicated system models that include and disregard friction can be evaluated with the direct method. The method is also useful in non-classical frameworks with similar properties, such as mass-energy conservation in special relativity theory. The equations characterizing systems in these and the classical framework are often too complex to solve analytically, and the direct method provides the only means by which attraction or stability domains of equilibria can be determined. In a wide variety of frameworks within physics, therefore, the direct method and concept of Lyapunov stability are indispensable.

4. Lyapunov Theory and Community Modeling. The brief outline of Lyapunov theory above suggests some advantages of defining ecological stability as Lyapunov stability. First, the definition formalizes the concept and integrates it into a well-known

mathematical theory. Stability properties of community models can then be assessed with analytical techniques like the indirect and direct methods.

Besides the obvious virtues of formalization, Lyapunov stability also seems to capture precisely the intuition that ecological stability depends upon community response to perturbation. Consulting (1), think of a community perturbed at time t_0 from \mathbf{x}^* to $\mathbf{x}(t_0)$ such that $|\mathbf{x}(t_0) - \mathbf{x}^*| < \delta$. If \mathbf{x}^* is Lyapunov stable, (1) states that the perturbed community will remain in an ε -neighborhood of \mathbf{x}^* for $\forall t \geq t_0$, *i.e.* the effects of the perturbation are circumscribed. Some other popular definitions of ecological stability are mathematically precise in that they are statistical functions of model variables (*e.g.* Lehman and Tilman's [2000] 'temporal stability'), but they do not, as Lyapunov stability does, provide a mathematical characterization of equilibrium system dynamics. If \mathbf{x}^* is Lyapunov stable in some region of E , for instance, the conditional in (1) ensures predictions can be made about system behavior after perturbation from \mathbf{x}^* . Mathematical definitions in the statistical sense cannot ground such predictions unless future values of variables depend upon past values in some systematic way, and the precise form of the dependency is known. The concept of Lyapunov stability is an important analytic tool within physics because (1) provides a mathematically precise description of the equilibrium dynamics of systems.

In addition, properties of community response to perturbation that are fundamental components of ecological stability *seem* to be formalizable with the direct method. The size of attraction domains of asymptotically Lyapunov stable equilibria and the rate systems return to them, for instance, are definable in terms of Lyapunov functions (Hahn 1963, §§12, 22-25). The larger the attraction domain, for instance, the

stronger the perturbations of variables a system can sustain and return to \mathbf{x}^* , which is often called system ‘tolerance’. Similarly, the “steeper” the Lyapunov function $V(\mathbf{x})$ (as gauged by $\nabla V(\mathbf{x})$), the faster it returns to \mathbf{x}^* , often called system ‘resilience’. If Lyapunov stability adequately defined ecological stability, it would therefore be possible to formalize the properties of tolerance and resilience and their relation to the stability of biological communities. Specifically, the intuitive idea that ecological stability increases with the strength of perturbation from which a system can return to equilibrium and with the speed of return to equilibrium after perturbation would be formalizable.

Finally, the direct method adequately characterizes the equilibrium dynamics of many systems in physics (and other sciences), and it is reasonable to expect the method will function similarly for ecological systems since their models share a similar mathematical structure. Like systems studied in physics, biological communities are usually modeled by differential equations. The only difference is that components of the vector $\mathbf{x}(t) = \langle x_1(t), \dots, x_i(t), \dots, x_n(t) \rangle$ represent biological variables, usually population sizes of species in a community, instead of typical physical variables.

This expectation seems to be supported, for instance, by the classical Lotka-Volterra model of one-predator, one-prey communities:

$$(6a) \quad \frac{dx_y(t)}{dt} = ax_y(t) - \alpha x_y(t)x_d(t);$$

$$(6b) \quad \frac{dx_d(t)}{dt} = -bx_d(t) + \beta x_d(t)x_y(t);$$

where x_d and x_y represent predator and prey population sizes; a represents prey birth rate; b represents predator death rate; and $\alpha, \beta > 0$ represent the effect of prey individuals on predator individuals and *vice versa*. Setting the left sides to 0 and solving yields one

nontrivial equilibrium \mathbf{x}^* where $x_d^* = \frac{b}{\beta}$ and $x_y^* = \frac{a}{\alpha}$. A Lyapunov function exists for all

$x_d, x_y > 0$:

$$(7) \quad V(\mathbf{x}) = c_1(x_y - x_y^* \ln x_y) + c_2(x_d - x_d^* \ln x_d);$$

where $c_1, c_2 > 0$ are constants, so \mathbf{x}^* is globally stable by the direct method (Logofet 1993).

An interesting property of (6) discovered by Vito Volterra reveals a structural similarity between models of ecological and physical systems, which in turn seems to justify the utility of Lyapunov theory in ecology. (6) can be rewritten as:

$$(8a) \quad \gamma_1 \dot{x}_y = \gamma_1 a x_y(t) - x_y(t) x_d(t);$$

$$(8b) \quad \gamma_2 \dot{x}_d = -\gamma_2 b x_d(t) + x_d(t) x_y(t);$$

where $\dot{x}_y = \frac{dx_y(t)}{dt}$, $\dot{x}_d = \frac{dx_d(t)}{dt}$, $\gamma_1 = \frac{1}{\alpha}$, and $\gamma_2 = \frac{1}{\beta}$. Adding these equations and

integrating yields:

$$(9) \quad \gamma_1 \dot{x}_y + \gamma_2 \dot{x}_d + \gamma_1 a \int_0^t x_y(t) - \gamma_2 b \int_0^t x_d(t) = c,$$

where c is a constant real value. Volterra believed (9) corresponded to a form of energy conservation [compare with (4) and (5) above].³ He called the first two terms “actual demographic energy,” and the second two “potential demographic energy” (Scudo 1971).⁴ If this structural similarity generalizes broadly across mathematical models of ecological systems, a view shared by another prominent founder of mathematical ecology Alfred Lotka (1956), it would seem to indicate that the stability definition used in physics is appropriate for ecology.

³ (9), in fact, is the underlying basis of (7). See Logofet (1993, 111-114).

⁴ At least for the first two terms, the resemblance with energy conservation is somewhat weak. The “actual demographic energy” corresponds closely with the mathematical form for the momentum of two particles, not their kinetic energy.

5. Lyapunov Stability in an Ecological Context. The adequacy of a definition depends upon what it requires of the defined concept. If ecological stability is defined as Lyapunov stability, for instance, this stipulates conditions biological communities must satisfy to be ecologically stable. If these conditions are unreasonable –if they are biologically unrealistic, too weak, or too strong for instance– the definition is inadequate. Despite its apparent advantages, interpreting Lyapunov stability in ecological terms furnishes strong reasons for rejecting the definition.

There are four types of Lyapunov stability to consider as definitional candidates for ecological stability. In order of increasing strength, they are local Lyapunov stability, nonasymptotic stability within a nonlocal domain, asymptotic stability within a nonlocal domain, and global asymptotic stability. Consider the weakest first. Although the frequent focus of community modeling, perhaps because it is relatively easy to evaluate with the indirect method (Hastings 1988), local stability defines ecological stability poorly. Differentiable models of systems can be linearized with the Jacobian matrix at any point of state space, such as at an equilibrium (see §3). By itself, however, linearization only provides information about local system behavior, *i.e.* only in an *infinitesimal* neighborhood of the linearization point. Local stability therefore says nothing about system behavior outside this extremely restricted domain. Besides it being fundamentally unclear how perturbation within an infinitesimal domain can be biologically interpreted (or empirically measured), real-world perturbations are clearly not of infinitesimal magnitude. Any real perturbation will expel a system at a strictly locally stable equilibrium from its stability domain. Local stability of nonlinear systems

like biological communities consequently provides no information about their response to common perturbations like drought, fire, population reduction by disease, *etc.* Application of the indirect method to community models, which can only evaluate local stability, therefore yields negligible insight into the community dynamics relevant to ecological stability.⁵

For different reasons, Mikkelson (1997) reaches a similar conclusion about local stability. His argument is based on the principle: “Do not employ a definition that turns what was originally thought to be an important empirical matter into a generic a priori exercise” (1997, 483). Applying this principle to the task of defining ecological stability, Mikkelson continues (1997, 485):

Requiring that every single species in a fifty-species community have a given property –*any* property– is a much stricter criterion than requiring that every species in a ten-species community have the same property. This means that as the number of species increases, the probability that the criteria are met almost certainly decreases. This leads to a *default* expectation that stability will decline with increasing diversity.

Since local stability requires *every* species return to its preperturbation value, Mikkelson claims that defining ecological stability as local stability decreases the likelihood species rich communities will be stable. It thereby biases *a priori* the debate about the relationship between community stability and diversity (construed as species richness) towards an inverse relationship, and violates his principle. On this basis, Mikkelson concludes the definition is indefensible.

The first problem with this argument is Mikkelson’s assumption that the likelihood of local stability of communities generally decreases with increasing species richness (1997, 494). This assumption is the underlying basis of the long quote above, but

⁵ Food web modeling of community stability is based on the indirect method, and has been criticized for this reason (Hastings 1988).

its truth is, however, an empirical rather than *a priori* issue. Interspecific interactions that emerge with greater species richness often increase the likelihood of stability, as the stabilizing effect of adding species to some community models shows (May 1974).

The second problem is that Mikkelson's principle is an unjustifiable adequacy condition for definitions in science because it confers too much import to unresolved scientific questions, regardless of how poorly formulated they may be. According to the principle, scientific concepts should not be defined in ways that resolve, even partially, outstanding questions that involve them. Yet scientific questions may themselves be ill-formed or confused due to ambiguity or vagueness of the concepts they concern. In fact, Shrader-Frechette and McCoy (1993) argue that the stability-diversity-complexity debate of community ecology is itself an example. Apparently intractable scientific debates and recalcitrant questions are sometimes justifiably resolved or, more accurately, dissolved when precise definitions of concepts reveal their misguided nature.⁶

The stronger concept of nonasymptotic Lyapunov stability within a nonlocal domain also inadequately defines ecological stability. Without asymptotic dynamics driving systems perturbed from equilibrium to return to it, successive weak perturbations can displace communities from their stability domains and, in the extreme case, cause species eradication. Ecological stability undeniably requires, however, persistence of the species of a community subject to successive weak perturbations.

Even with asymptotic dynamics, however, Lyapunov stability does not define ecological stability adequately. To see why, consider defining ecological stability as the strongest Lyapunov stability property: global asymptotic stability. Unlike the Lyapunov

⁶ Einstein's definition of simultaneity, for instance, arguably settled a number of scientific questions about temporal relations between events.

stability properties considered thus far, global asymptotic stability appears to be much stronger than ecological stability because it seems to entail that a community will return to equilibrium following any perturbation that does not eradicate its species. Model variables of systems at asymptotically and globally stable equilibria that are perturbed to 1% of their equilibrium values, for instance, will deterministically return to their initial values. This response to such severe perturbations is not required for a biological community to be ecologically stable, but it certainly seems sufficient.

Despite its apparent plausibility, the idea that global asymptotic Lyapunov stability is sufficient (but not necessary) for ecological stability is unjustified. It is based on a flawed conception of ecological perturbation. Real-world perturbations to biological communities do not merely affect population sizes, represented by variables in models like (2). They also change environmental factors that influence community dynamics, represented by model parameters. As May (1974, 216) creatively put it:

in nature, population perturbations are driven not by the stroke of a mathematician's pen resetting initial conditions [*i.e.* resetting variable values to represent perturbations], but by fluctuations and changes in environmental parameters such as birth rates, carrying capacities, and so on.

Perturbations may induce, for instance, changes in the strength and qualitative nature of interspecific and intraspecific community interactions. Initially noninteracting species may begin competing or exhibiting other non-neutral interactions, and *vice versa*. Since real-world perturbations may affect environmental factors that influence community dynamics as well as population sizes of species, they should be represented by corresponding changes in variable *and* parameter values. Ecological stability should then be evaluated with respect to system behavior following both types of changes.

Lyapunov stability, however, only assesses system response to temporary changes in *variable* values (see [1] above). It does not consider system responses to changes in parameters and thus provides only a partial account of the type of system response to perturbation ecological stability requires. Lyapunov's direct method is therefore not a suitable methodology for analyzing ecologically relevant nonlocal stability properties of standard community models, although it accurately estimates stability to perturbations affecting only model variables (see below).

This difference between Lyapunov and ecological stability consequently leads to different evaluations of the stability of mathematical models of biological communities. Consider the generalization of (6) to n -species:

$$(10) \quad \frac{dx_i}{dt} = x_i \left(r_i - \sum_{j=1}^n \gamma_{ij} x_j \right);$$

where r_i designates the intrinsic growth rate of species i ; γ_{ij} designates the effect of species j on species i ; and $1 \leq i, j \leq n$. (10) can represent several realistic features of biological communities. The γ_{ii} designate intraspecific interactions and the γ_{ij} ($i \neq j$) can represent any type of interspecific interaction: predator-prey $\gamma_{ij}\gamma_{ji} < 0$, competition $\gamma_{ij}, \gamma_{ji} < 0$, mutualism $\gamma_{ij}, \gamma_{ji} > 0$, etc.

Two types of systems, called "conservative" and "dissipative," have been studied with (10) and highlight the difference between ecological and Lyapunov stability particularly clearly. If:

$$(11) \quad \sum_{i,j=1}^n \gamma_{ij} x_i x_j \geq 0,$$

the system is dissipative with strict inequality, and conservative with equality. Informally, (11) says interspecific interactions do not influence total system dynamics in conservative

communities and retard them in dissipative communities. Dissipative and conservative communities, Logofet (1993) suggests, are analogous to mechanical systems with and without friction.

In conservative (dissipative) communities: (i) $\gamma_{ii}=0$ ($\gamma_{ii}>0$) for all i ; and, (ii) $\gamma_{ij}=-\gamma_{ji}$ for all $i\neq j$. (i) entails there is intraspecific self-damping in dissipative communities, and none in conservative communities. (ii) entails the communities only exhibit antisymmetric interspecific predator-prey interactions. Note that (10) is not necessarily a two trophic level model of $\frac{n}{2}$ predators and $\frac{n}{2}$ prey. Species i may be the predator and prey of different species, and part of k -length ($k\leq n$) food chains.

The direct method shows \mathbf{x}^* is a nontrivial equilibrium, *i.e.* $(\forall i)(x_i^* > 0)$, of conservative n -species community models iff \mathbf{x}^* is globally Lyapunov stable; it is asymptotically and globally Lyapunov stable iff the community model is dissipative. These results only hold, however, if n is even. As an indication of what defining ecological stability as global Lyapunov stability would require of communities modeled by (10) and (11), this restriction seems arbitrary and counterintuitive. Without any empirical evidence or a biological reason to suppose there is a relationship between evenness or oddness of species richness and ecological stability, of which there seems to be none, this requirement is unreasonable.⁷

A second, more revealing feature of these results is their dependence on the exact anti-symmetry of $[\gamma_{ij}]$. If jeopardized even slightly, they no longer hold and the system may become unstable (Levin 1981). Small changes in the values of these parameters

⁷ Of course, it is possible such a relationship exists, but has not yet been found. Most ecologists highly doubt this claim. May (1974, 53), for instance, suggested it, “border[s] on the ridiculous,” and Volterra, who first recognized the fact, was similarly skeptical (Scudo 1971).

caused by realistic but weak perturbations may therefore undermine asymptotic global stability and initiate instability. This occurs because Lyapunov stability defines stability strictly in terms of system behavior after changes to variables, not after changes to parameters as well. For this reason, asymptotic global stability and weaker Lyapunov stability concepts poorly define ecological stability.

The disparity between what the direct method says about the stability properties of biological communities and what ecological stability seems to require is not limited to conservative and dissipative models of communities. A wide range of community models the direct method shows are asymptotically stable in nonlocal domains are similarly sensitive to slight changes in parameters (see Hallam 1986 and references therein).

In general, therefore, Lyapunov stability inadequately captures the intuition that ecological stability depends on community response to perturbation because it inadequately represents real-world perturbations. As such, Lyapunov stability fails to capture principal features of ecological stability like resilience and tolerance. With respect to abiotic perturbations such as temporary temperature or precipitation anomalies, for instance, resilience and tolerance cannot be measured using the direct (or indirect) method (see §4). Temporary changes in model parameters representing these environmental factors, not changes to model variables, best represent these types of perturbation. Resilience and tolerance should be gauged by the subsequent changes in variables following these changes. Specifically, in more resilient systems, variables return more rapidly to their preperturbation values following temporary changes in parameters. Similarly, the greater the parameter change the system can sustain and have its variables return to equilibrium, the more tolerant it is.

Lyapunov stability, however, cannot represent these features of ecological stability properly because it cannot accurately represent community response to perturbations that affect factors represented by model parameters rather than just variables. Since, tolerance and resilience of biological communities depend upon how communities respond to perturbations that affect model variables *and* parameters, they therefore cannot be defined in terms of properties of Lyapunov functions (see §3). Whereas Lyapunov stability focuses strictly on system behavior in *state* space, ecological stability essentially concerns system behavior in *parameter* space as well. Consequently, Lyapunov stability is not a sufficient condition for ecological stability, as some ecologists have explicitly suggested (*e.g.* Logofet 1993, 109) and many ecologists assume.

6. Conclusion. That ecological stability should not be defined as Lyapunov stability does not entail the latter should play no role in mathematical modeling of biological communities. For a community at equilibrium, asymptotic Lyapunov stability within nonlocal attraction domains adequately represents ecologically stable responses to perturbations that alter population sizes of community species but leave community structure and parameter values unchanged. Lyapunov stability is therefore a plausible necessary condition for ecological stability. Moderate culling of some community species or temporary cessation of plant or animal harvesting in ecosystems where it has a long history are plausible examples of such perturbations. In fact, the effects fishing cessation during WWI would have on fish populations motivated Volterra to begin developing mathematical models of communities (Scudo 1971).

This analysis also does not detract from the obvious gains in precision, rigor, and other scientific virtues afforded by mathematical dynamic systems modeling within ecology. It does expose, however, a limitation of the stability definition that usually accompanies this type of modeling. If ecological systems like communities were completely analogous to systems studied within physics for which Lyapunov's methods are so successfully applied, defining ecological stability as Lyapunov stability would be entirely justifiable. Section 5 makes clear there is, however, a crucial difference in how perturbations should be represented in the two contexts. Scientific definitions that make an initially problematic concept precise by integrating it into a systematically developed mathematical theory, even one fruitfully utilized in other sciences, are therefore not always satisfactory.

Within classical mechanics, force fields governing interactions between bodies, such as gravitation, are usually invariant. Masses of bodies in the system, overall system structure (*e.g.* mass-spring system, damped pendulum, mass-pulley system, *etc.*), and other system features are also usually held or assumed fixed in this framework. Model features representing these system properties, such as parameter values and model structure, are correspondingly fixed. With the background held constant in this way, perturbations are represented by temporary finite changes in variable values, for instance, by displacements of position vectors and alterations to velocity vectors (see §3). Energy is conserved or monotonically dissipated in these systems depending upon whether they are closed or open, so Lyapunov functions can always be constructed for them and their stability properties evaluated. Since perturbations of such systems are not taken to change

the system properties represented by parameters and model structure, Lyapunov stability captures their important stability properties.

Within ecology, however, parameters representing external factors affecting species and their interactions are much more likely to change than in systems studied within classical mechanics because they are regularly altered by real-world perturbations.⁸ An appropriate definition of ecological stability therefore requires integrating Lyapunov stability with a concept representing how communities respond to these types of structural change: a structural stability concept. This would adequately account for how communities respond to real-world perturbations represented by temporary changes in variable *and* parameter values. Resilience and tolerance to real-world perturbations would be representable with such a concept.

This suggestion poses a formidable challenge to mathematical ecologists because structural stability is a much more technically complex mathematical concept than Lyapunov stability (see Peixoto 1959). The dearth of work, especially biologically-oriented work, devoted to it (Lewontin 1969; May 1974), however, may explain its lack of application within mathematical ecology, not any essential mathematical intractability of the concept.

The first prominent mathematical ecologists, Lotka and Volterra, were physicists by training and this significantly influenced their approach to modeling biological systems (Kingsland 1995). Their work, moreover, subsequently set much of the agenda of twentieth century mathematical ecology. Not surprisingly, most mathematical

⁸ The difference between ecological systems and those studied in physics generally should not be overstated, however. In fluid mechanics, for instance, the background structure of the systems studied is highly variable. For the reasons discussed in section 5, therefore, Lyapunov stability would not be an appropriate representation of the stability of these systems.

ecologists have used the concept of Lyapunov stability and the direct and indirect methods to analyze community models. A stability concept that incorporates structural stability, however, constitutes a more defensible definition of ecological stability, and would better characterize the dynamics of ecologically stable biological communities.

References

- Goh, B. S. (1977), "Global Stability in Many-Species Systems." *American Naturalist* **111**: 135-143.
- Hahn, Wolfgang (1963), *Theory and Application of Liapunov's Direct Method*. Englewood Cliffs: Prentice-Hall.
- Hallam, Thomas (1986), "Population Dynamics in a Homogeneous Environment." In Hallam, Thomas and Levin, Simon (eds.) *Mathematical Ecology: An Introduction*. New York: Springer, 61-94.
- Hastings, Alan (1988), "Food Web Theory and Stability." *Ecology* **69**: 1665-1668.
- Hinrichsen, Diederich and Pritchard, Anthony (2005), *Mathematical Systems Theory I*. New York: Springer.
- Kingsland, Sharon (1995), *Modeling Nature*. 2nd edition. Chicago: University of Chicago Press.
- Lehman, Clarence and Tilman, David (2000), "Biodiversity, Stability, and Productivity in Competitive Communities." *American Naturalist* **156**: 534-552.
- Levin, Simon (1981), "The Role of Mathematics in Biology." *Proceedings of the Landmoedt om Matematikken I*. Copenhagen: Danish Mathematical Society pp. 455-478.
- Lewontin, Richard (1969), "The Meaning of Stability." In Woodwell, G. and Smith, H. (eds.) *Diversity and Stability in Ecological Systems*. Brookhaven, NY: Brookhaven Laboratory Publication No. 22,13-24.
- Lotka, Alfred (1956), *Elements of Physical Biology*. New York, NY: Dover Publications.
- Logofet, Dmitrii (1993), *Matrices and Graphs: Stability Problems in Mathematical Ecology*. Ann Arbor: CRC Press.
- Lyapunov, Aleksandr ([1892], 1992), *The General Problem of the Stability of Motion*. London: Taylor and Francis.
- May, Robert (1974), *Stability and Complexity in Model Ecosystems*. 2nd Edition. Princeton: Princeton University Press.
- Mikkelsen, Greg (1997), "Methods and Metaphors in Community Ecology: The Problem of Defining Stability." *Perspectives on Science* **5**: 481-498.

Odenbaugh, Jay (2001), "Ecological Stability, Model Building, and Environmental Policy: A Reply to Some of the Pessimism." *Philosophy of Science (Proceedings)* **68**: S493-S505.

Peixoto, M. M. (1959), "On Structural Stability." *Annals of Mathematics* **69**: 199-222.

Scudo, Francesco (1971), "Vito Volterra and Theoretical Ecology." *Theoretical Population Biology* **2**: 1-23.

Shrader-Frechette, Kristin and McCoy, Earl (1993), *Method in Ecology*. Cambridge, UK: Cambridge University Press.

Figure 1. Graphical representation of nonasymptotic and asymptotic Lyapunov stability.

