

How to Fix Kind Membership:

A Problem for HPC-Theory and a Solution

Abstract

Natural kinds are often contrasted with other kinds of scientific kinds, especially functional kinds, because of a presumed categorical difference in explanatory value: natural kinds can ground explanations, while other kinds of kinds cannot. I argue against this view of natural kinds by examining a particular type of explanation – mechanistic explanations – and showing that functional kinds do the same work there as natural kinds are supposed to do in more standard scientific explanations. Breaking down the categorical distinction between natural kinds and other kinds of kinds, I argue, delivers two goods: It provides us with a view of natural kindhood that does justice to the epistemic roles of kinds in scientific explanations. And it allows us to solve a problem that HPC-theory, currently one of the most popular accounts of natural kindhood, confronts.

Word count: 4972 words, counting main text, title, abstract, footnotes and references.

1. Introduction. What should any good theory of natural kindhood minimally provide us with? While different philosophers entertain different views of what the natural kinds debate is actually about and what exactly a theory of natural kinds should deliver, two demands seem beyond doubt. First, any philosophical theory of natural kinds should specify what distinguishes natural kinds from other kinds of kinds. Second, any natural kind theory should specify what sorts of factors determine the kind membership of any given entity.

In the context of philosophy of science, these demands take on a specific form. Natural kinds are often contrasted with other kinds of kinds featured in science, in particular functional kinds, on the grounds of a presumed categorical difference in explanatory value. Natural kinds are epistemologically privileged, the thought is, because reference to them has explanatory force, whereas reference to other kinds of kinds does not. And the underlying reason why natural kinds can ground explanations is that they (in contrast to other kinds of kinds) group things according to properties on which scientific explanations zoom in, such as the “real natures” of things (as typically claimed in essentialist accounts of natural kindhood) or those properties that appear in laws of nature (e.g., Fodor, 1974; Churchland, 1985). Accordingly, scientific fields that do not focus on natural kinds but classify their subject matter into “mere” functional and other kinds of kinds – i.e., virtually all of the special sciences – are often deemed non-explanatory and hence non-autonomous, non-fundamental sciences.

My aim in the present paper is to counter this widespread intuition that natural kinds are epistemologically “special”. In so doing, I aim to show how one of the currently most popular accounts of natural kindhood, Richard Boyd’s “homeostatic property cluster”

(HPC) theory, can be modified to accommodate the view of natural kinds that I defend. My strategy will be to focus on the standard contrast class to the category of natural kinds, namely functional kinds, and to proceed as follows. First, I shall point to a problem that HPC-theory confronts (Section 2). I shall continue in Section 3 by examining how functional kinds are featured in a particular type of scientific explanation, namely mechanistic explanation (henceforth: ME). If functional kinds do the same work in MEs as accepted natural kinds do in “standard” explanations and if MEs can be counted as genuinely explanatory, then functional and natural kinds should be placed on an equal footing with respect to their epistemological importance. I shall argue that the two antecedents of this argument are indeed the case (Sections 3.1 and 3.2, respectively) and, therefore, natural kinds and the sciences that focus on them are not epistemologically “special”. I shall conclude in Section 4 by bringing the two lines of work from Sections 2 and 3 together: the realization that there is no in principle difference between functional and natural kinds regarding their epistemic importance for scientific reasoning and explanation provides clues how the problem for HPC-theory can be resolved and HPC-theory can be turned into a full-fledged theory of natural kindhood able to cover kinds in the special sciences as well as the less controversial natural kinds.

2. A Problem for HPC-theory. In the philosophical literature on the topic two distinct ways of thinking about natural kinds can be found. On the one hand there is the essentialist tradition that, broadly taken, understands natural kinds as grouping of things according to their natures, their intrinsic properties or causal capacities, their microscopic structures, etc. On the other hand there is the more recent tradition that understands

natural kinds as groupings of things over which we can make reliable inductions. That these lines of work really are quite distinct can be seen from the way in which they conceive of the problem of natural kinds.¹ The former line of work conceives of the problem as a metaphysical problem, i.e., as the question what sorts of things there are in the world. As Brian Ellis put it in a recent defense of essentialism, “membership of a natural kind is decided by nature, not by us” (2001, 19). The latter line of work, in contrast, sees it as foremost a question in epistemology, i.e., as the question what ways of grouping things help us to make inferences and to explain phenomena. Boyd, for example, asserted that “[i]t is a truism that the philosophical theory of *natural* kinds is about how classificatory schemes come to contribute to the epistemic reliability of inductive and explanatory practices” (1999b, 146; also 1999a, 69; the emphasis is Boyd’s). On this view, kind membership is decided more by us than by nature.

With respect to the explanation-grounding capacity of natural kinds, the two lines of work provide different answers that run into different kinds of problems. From the perspective of the essentialist tradition it should be no miracle that natural kinds ground reliable inferences and explanations. If there is a definitive, objective account of what kinds of things make up the furniture of the world, then clearly any explanation of a given phenomenon should ultimately make reference to the kinds of things that are involved in the phenomenon under consideration. The problem, however, is that we do not have any direct access to the world that would allow us to compile the required definitive inventory of the world’s furniture. Our best bet at obtaining such an inventory

¹ The distinction is found (albeit implicitly) in a discussion between Hacking (1991) and Boyd (1991). Neither of these two authors, however, has actually claimed that there are two distinct traditions of thinking about natural kinds.

is to consult the various fields of science and to look at the ontologies that these currently adopt. But scientists entertain particular ontologies because these make sense in the context of particular theories – more specifically, because the kinds included in these ontologies can serve as the bases of generalizations, explanations, predictions, etc. *against the background* of the particular theories that are adopted. And this brings us back to the question that we started with, i.e., wherein lies the explanation-grounding capacity of the kinds that particular theories recognize.

The second, epistemology-oriented tradition of thinking about natural kinds thus seems to be in a better position than the essentialist tradition, as it begins by looking at how reference to kinds is epistemically important in actual scientific practice. Boyd's HPC-theory is, as indicated above, at present the most prominent exponent of this line of work.² It was developed from the 1980s onward as an attempt to take seriously the epistemic roles that kinds play in the special sciences, regardless of whether or not they fit the essentialist picture of natural kinds. HPC-theory starts from the recognition that most kinds that feature in science are not groupings of things with *exactly* the same (microstructural, causal, ...) properties, but groups of things that bear various degrees of causally supported resemblances to each other, that is, they all exhibit largely similar properties due to largely the same causes (Boyd, 1999b, 142-144). Accordingly, kinds should not be defined by separately necessary and jointly sufficient essential properties that all and only the members of the kind exhibit without exception, but by the cluster of properties that are found to regularly, but not exceptionlessly, cluster together in natural

² Primers of HPC-theory can be found in Boyd (1999a; 1999b).

entities in combination with the set of causal factors (Boyd speaks of “homeostatic mechanisms”) that underlie this clustering of properties:³

(HPC) a particular natural kind is defined by a combination of a particular F and a particular H , where

F = the set of all properties that are found to repeatedly cluster in nature, where this clustering may be imperfect and exception-ridden (the “property family”),

H = the set of causal factors (“homeostatic mechanisms”) that underwrite this clustering.

Because for a given natural kind there is no set of properties unique to the members of that kind, F cannot exhaustively define the kind. Accordingly, HPC-theory adds H to the definition and assumes the combination of F and H to uniquely define a kind: a kind is defined by the properties that are found to repeatedly cluster together plus the underlying factors that cause this clustering.

In order to do justice to the messy state of affairs in the world in which entities are hardly ever exactly alike, HPC-theory conceives of the F and H that define a particular natural kind in an *open-ended* manner: no property is necessarily unique to one F , no causal factor is necessarily unique to one H , the F of a kind may come to include new properties and present properties may cease to be members, causal factors may begin or cease to operate, and there are no “core sets” of properties or underlying causal factors that all and only members of the corresponding kind exhibit or are affected by. This

³ The notation in terms of F and H is mine.

yields an account of natural kinds that is sufficiently flexible to accommodate all kinds that feature in the various special sciences, as well as the traditional natural kinds.

However, precisely this flexibility causes a problem for HPC-theory. Essentialist accounts of natural kinds tell us which factors in nature determine the extensions of kinds. If for a particular kind a kind essence is identified, we immediately have a criterion for assessing whether or not any given entity is a member of that kind: Does it instantiate the kind's essence; does it exhibit all the properties deemed necessary and sufficient for membership in the kind? (This may often not be a very operational criterion, but it at least is a criterion that can be used in principle.) HPC-theory, however, fails to provide us with any such criteria. Even if we have fully identified all the members of the property family F and of the set of causal factors H for a given kind, we still have no criteria for determining the kind's extension. The reason is that both F and H are open-ended in the abovementioned sense. Both the F and H that define a particular kind may *in principle* change in time to such an extent that at a later time they no longer contain any of the elements that they contained at an earlier time.

This can be seen particularly clearly in the case of biological species that Boyd presented as prime examples of HPC-kinds (Boyd, 1991; 1999a; 1999b). According to Boyd, biological species are HPC natural kinds defined by the properties that a species' member organisms typically exhibit in combination with the mechanisms that underwrite this clustering (i.e., common descent and reproductive cohesion – Boyd 1999b, 167). But species are subject to open-ended evolution: its organisms can come to exhibit newly evolved properties and old traits can be lost as time goes by, while there is no reason to assume that any particular core set of properties will be conserved. Furthermore, in the

case of a speciation event in which a new species branches off from its ancestor species, the member organisms of the two species will typically be characterized by the same family of properties. The same holds for H if the relevant causal factors are, for example, environmental: the environment may change heavily during a species' lifetime or remain the same over the lifetime of an ancestral species and a series of its descendants. Hence, the combination of F and H is insufficient to determine the boundaries of the species that it is supposed to define.

HPC-theory, then, can only account for kinds the extensions of which have already been fixed independently by other means. If we have independent criteria by which we can enumerate exactly which entities are to be counted as the members of a given kind and which are to be discounted, HPC-theory can tell us which properties and underlying causal factors hold the kind together. (In the case of species, the relevant independent criterion is organisms' locations on particular branches of the Tree of Life.) But a good natural kind theory should do more: it should also provide us with criteria with which kind membership can be determined in the first place – and on this count HPC-theory fails.

3. Functional Kinds in Explanations. Clues about how to solve this problem for HPC-theory can be obtained from an examination how kind terms function in actual scientific explanations. In “standard” explanations the names of natural kinds typically appear – if not in statements of laws of nature – in statements of lawlike generalizations of the form

“All *Ks* have property *p*” that can ground explanations.⁴ But at the same time the natural kinds that are being referred to are themselves phenomena in need of an explanation: why do elementary particles, atoms, etc. come in those kinds that they do, rather than different ones? Reference to natural kinds, then, has a double role in scientific explanation: natural kinds are mentioned as explananda and as explanantia.

In what follows, I shall argue that this double role is not unique for natural kind names. In some types of explanations functional kinds play the same epistemic roles as “standard” natural kinds play in “standard” explanations. Clearly, a complete analysis of the explanatory value of reference to kinds cannot be undertaken here, as the variety of classificatory and explanatory strategies used in the various sciences is just too large. I shall therefore argue by example and focus on one particular type of explanation that is increasingly moving into the focus of philosophy of science, namely mechanistic explanation.

3.1. Mechanistic explanation. At present, MEs are increasingly receiving philosophical attention in what has come to be called the “new mechanistic philosophy”.⁵ Although the various authors in the “new mechanistic philosophy” advocate different accounts of how MEs explain, they share the basic view that MEs explain “by describing how the component entities and activities are organized together such that the phenomenon

⁴ The often assumed connection between laws and kinds is not straightforward, as the paradigmatic laws of nature do not typically mention natural kinds. But this is an issue that I must leave for elsewhere.

⁵ For brief overviews of the “new mechanistic philosophy”, see Skipper and Millstein (2005) or Craver (2006).

occurs” (Craver 2006, 374). That is, an ME of a particular property or behavior of a given entity proceeds by (1) decomposing the entity under study into its constituent parts, each of which exhibits a particular behavior when placed in a particular systemic context, and (2) specifying the actual systemic context in which each of the parts is embedded. The explanandum then can be derived from generalizations about the characteristic behavior of the constituent parts combined with specifications of the actual systemic conditions.

Examples of MEs are found throughout the various special sciences. Let me here consider an ME from the domain of evolutionary-developmental biology in some more detail: the *Drosophila* Segment Polarity Network (SPN) (Von Dassow and Munro 1999; Von Dassow et al. 2000). The SPN is a gene regulatory network consisting of five genes (named “*engrailed*”, “*hedgehog*”, “*wingless*”, “*cubitus interruptus*” and “*patched*”) and responsible for inducing a head – rear end orientation in fruit fly embryos. The developmental explanandum – in this case the appearance of polarity in the body segments of developing *Drosophila* embryos – is explained by reference to, among other things, the typical behavior of SPNs in the context of a particular stadium of embryo development. The particular behavior of SPNs, in turn, is taken as explanandum and is explained by reference to the typical behavior of the various kinds of entities of which SPNs are composed and the way in which these interact in the context of the SPN. Here, ‘*engrailed*’, ‘*hedgehog*’, etc. appear as functional kind names, as genes are identified by their functions – that is, causal roles – in molecular studies (e.g., Waters 1994; Griffiths and Stotz 2007). In a next step, the particular behavior of entities of, say, the *engrailed* kind can be taken as the explanandum and explained by reference to the various kinds of

functional parts of which *engrailed* genes are composed (*cis*-regulatory modules, exons, etc.).

This explanatory strategy is recursive. In each step, part of the explanation is achieved by specifying the typical intrinsic behavior of a system's constituent parts while black-boxing these parts themselves. What matters in MEs is that particular parts perform particular causal roles under particular circumstances. How these causal roles are actually realized is not important in the analysis of the overarching system – it becomes interesting only when the functional part is considered in isolation and the research question becomes which behaviors it exhibits in which circumstances. Functional kinds, then, serve as the “hinges” around which MEs turn in the following sense. Reference to the functional kind to which a particular part of a system belongs is explanatory as the basis of a generalization about the behavior that it is expected to exhibit when placed in a particular environment. In addition, the existence of the various functional kinds is itself a phenomenon in need of an explanation, as it needs to be explained how the black-boxed entities are able to realize the various functions that they realize in different systems. Thus, functional kinds play a double role in MEs as explanantia and the explananda in the same way as commonly accepted natural kinds play a double epistemic role in more “standard” scientific explanations. Just as natural kinds can be said to be the “hinges” around which “standard” scientific explanations turn, functional kinds can be conceived of as the “hinges” of MEs.

3.2 Are MEs genuinely explanatory? What I have said, however, does not imply that reference to functional kinds in MEs is actually explanatory in the same sense as citing

natural kinds is. The fact that a system like the *Drosophila* SPN can be hierarchically decomposed into a number of subsystems does not imply that such a decomposition has any explanatory force. After all, any material entity can be decomposed into parts in a plethora of different ways, without all these possible decompositions necessarily picking out kinds with the same degree of explanatory import or even any explanatory import. What makes reference to natural kinds explanatory is that natural kinds are supposed to represent objectively existing features of nature, rather than mere ways of grouping things useful for our particular purposes. Similarly, I want to suggest, MEs can be considered genuinely explanatory when the decomposition of the system under consideration into functional parts can be understood as representing not just a heuristically useful way of analyzing a given system, but as identifying organizational structures that are actually found in nature.

In the case of systems like the *Drosophila* SPN, this suggestion can be clarified by taking recourse to the notion of modularity. Roughly, modularly organized systems are systems that are not composed of their basic parts in a simple, aggregative manner (like bricks stacked in a wall), but exhibit a multi-level compositional structure (a system composed of functionally interdependent subsystems, in turn composed of functionally interdependent sub-subsystems, in turn composed of sub-sub-subsystems, etc. until the level of the basic parts is reached). Each of these units and subunits – the *modules* into which the system can be decomposed – are comparatively well-integrated subsystems of the larger system that are (to a good degree) materially separable from other subsystems of the same system, built of various recognizable components and distinguishable from other modules by the well-defined functions (causal roles) that they perform in the

context of the system of which they are parts (Von Dassow & Munro 1999, 307; Bolker 2000). Although modularity has been at the focus of scientific and philosophical attention for some time now, it has turned out surprisingly difficult to define precisely what modules are (e.g., Von Dassow & Munro 1999, 312; Bolker 2000, 771; Rieppel 2005, 18). For my purposes, however, it is sufficient to notice that scientists have good reasons to conceive of modules as objective features of natural systems.

Over 40 years ago, Herbert Simon (1962) pointed out that we can expect to find modularly organized systems in the living world, because of two ways in which organizing systems in a modular way can be advantageous. Modular organization can contribute to the efficiency of the assembly processes in which systems come into being and it can contribute to the functional stability of finished systems. That is, modular organization can positively contribute to what is commonly called the “evolvability” of biological systems (their capacity to evolve further) and to their “survivability” (their ability to survive in various environments).⁶ At present, it is increasingly becoming clear that modularity is indeed a widely found property of systems in the living world and that various sorts of modules can be identified on many different levels of organization (Bolker 2000, 774; Callebaut 2005).

There is, then, no reason to conceive of the various functional kinds of modules that are featured in MEs of systems like the *Drosophila* SPN as being less representative of objective features of nature than “standard” natural kinds. Therefore, *if* one is prepared to admit that commonly accepted natural kinds play important epistemic roles in “standard”

⁶ For a discussion of the notion of evolvability in recent biological literature, see Love (2003). ‘Survivability’ is my own term.

scientific explanations (something that I want to leave open for the moment, however), then one should also be prepared to admit that *some* functional kinds play the same roles in MEs.

4. Fixing HPC-theory. If I am correct in my suggestion that there is no in principle difference between functional and natural kinds regarding their epistemic importance for scientific reasoning and explanation, this could help to remedy the abovementioned problem for HPC-theory in the following manner. The root of the problem, I believe, is that, even though Boyd repeatedly emphasized that kinds are dependent on the disciplinary context in which they feature, HPC-theory still rests on too realist a view of natural kinds. Boyd (1991, 141-142) himself pointed to the fact that the HPC-definition of a kind often fails to fully specify kind membership, but did not consider this a problem for HPC-theory. According to Boyd, this indeterminacy is a necessary element of the HPC-definition of kinds, as it reflects the actual state of affairs in nature in which many kinds are a bit “vague around the edges”. Boyd’s phrasing often suggests that stable property clustering is the normal case found “out there” in the world and exceptions, while occurring regularly, still constitute the minority of cases.

But, as the case of MEs shows, there are scientific explanations that derive explanatory force from kinds that are not defined by families of properties found to cluster repeatedly in natural entities. MEs use functional kinds in which kind membership is determined by the capability of performing a particular causal role in the context of a particular system, irrespective of the particular properties of the entities that perform these functions. The particular properties relevant for performing the function under

consideration only come into focus when the functional kind itself is taken as explanandum. Taking seriously Boyd's suggestion that "the philosophical theory of *natural* kinds is about how classificatory schemes ... contribute to ... inductive and explanatory practices" (1999b, 146; quoted above) implies that we should focus on those criteria that scientists *actually* use when identifying kinds for use in explanations. When these criteria are not framed in terms of properties that the kind's members exhibit, we should turn to those factors that actually make the kind's members explanatorily interesting (for instance a particular function). This suggests the following modification to (HPC) as presented in Section 2:

(HPC*) a particular natural kind is defined by a combination of a particular Φ , F^* and H^* , where

Φ = the factor that makes members of the kind explanatorily interesting (e.g. the capability to perform a particular causal role function),

F^* = the set of properties that play central roles in the explanation of Φ and are found to repeatedly cluster in nature, where this clustering may be imperfect and exception-ridden,

H^* = the set of causal factors that underwrite this clustering.

The definition of a natural kind that is obtained in this way does justice to the following important motivation behind Boyd's account.⁷ Natural kinds do not simply emerge from a direct examination of the state of affairs in nature, nor do they emerge

⁷ A motivation that has also been put forward earlier, for example by Platts (1983).

exclusively on the basis of whatever way of classifying things we might find useful in particular contexts. Rather, natural kinds emerge from human interactions with nature in investigative and explanatory practices. In many cases, the kinds that emerge there cannot be defined by those explanatorily important properties that repeatedly cluster together in natural entities, as these properties are not specified in the explanations under consideration. This was the case in the MEs discussed in the previous section, in which the explanatorily important characteristics were functional and the corresponding explanatory kinds were functional kinds.

Summarizing, my suggestion how to fix HPC-theory is as follows. Traditional accounts of natural kinds conceive of kinds as identified by clusters of metaphysically singled-out properties that constitute the essences of the kinds' members. HPC-theory counters that the relevant properties are not metaphysically but epistemically singled out: they are explanatorily important properties and hence crucially depend on the investigative context in which a kind is used. But as there are cases in which the explanatorily important properties are left unspecified in explanatory relevant kinds, defining kinds in terms of property clustering is not always adequate to the investigative or explanatory practice under consideration. This can be remedied by adding the factor that *actually* makes a kind explanatorily interesting to the definition of the kind.

5. Concluding remark. I have argued that functional kinds should not be thought of as *intrinsically* explanatorily unimportant or even less important than “good” natural kinds. As functional kinds do the same work in MEs as accepted natural kinds do in “standard” scientific explanations *and* MEs are genuinely explanatory, the functional kinds that

feature in MEs should not be conceived of as being categorically distinct from natural kinds. But it is important to see that I have not advanced a general argument that functionally defined kinds should without further ado be brought under the natural kind fold, or for a suggestion that modified HPC-theory could be used to do this. Many functional kinds don't have much explanatory power, but *some* do and should therefore be thought of as natural kinds. I have suggested a modification to Boyd's account of natural kinds that takes seriously the suggestion that natural kinds are kinds of things that stand at the focal points of scientific explanations. Focusing explicitly on the explanatory role of kinds in science yields, I believe, an account of natural kinds that is more appropriate to those kinds that are featured in the various special sciences than is the original formulation of HPC-theory.

References

- Bolker, Jessica A. (2000), "Modularity in Development and Why it Matters to Evo-Devo", *American Zoologist* 40: 770-776.
- Boyd, Richard N. (1991), "Realism, Anti-Foundationalism and the Enthusiasm for Natural Kinds", *Philosophical Studies* 61: 127-148.
- (1999a), "Kinds, Complexity and Multiple Realization", *Philosophical Studies* 95: 67-98.
- (1999b), "Homeostasis, Species, and Higher Taxa", in Robert A. Wilson (ed.), *Species: New Interdisciplinary Essays*. Cambridge (Mass.): MIT Press, 141-185.

- Callebaut, Werner (2005), "The Ubiquity of Modularity", in Werner Callebaut and Diego Rasskin-Gutman (eds), *Modularity: Understanding the Development and Evolution of Natural Complex Systems*. Cambridge (Mass.): MIT Press, 3-28.
- Churchland, Paul (1985), "Conceptual Progress and Word/World Relations: In Search of the Essence of Natural Kinds", *Canadian Journal of Philosophy* 15: 1-17.
- Craver, Carl (2006), "When Mechanistic Models Explain", *Synthese* 153: 355-376.
- Ellis, Brian (2001), *Scientific Essentialism*, Cambridge: Cambridge University Press.
- Fodor, Jerry A. (1974), "Special Sciences (Or: The Disunity of Science as a Working Hypothesis)", *Synthese* 28: 97-115.
- Griffiths, Paul E. and Karola Stotz (2007), "Gene", in David L. Hull and Michael Ruse (eds), *The Cambridge Companion to the Philosophy of Biology*, Cambridge: Cambridge University Press, 85-102.
- Hacking, Ian (1991), "A Tradition of Natural Kinds", *Philosophical Studies* 61: 109-126.
- Love, Alan C. (2003), "Evolvability, Dispositions, and Intrinsicity", *Philosophy of Science* 70: 1015-1027.
- Platts, Mark (1983), "Explanatory kinds", *British Journal for the Philosophy of Science* 34: 133-148.
- Rieppel, Olivier (2005), "Modules, Kinds, and Homology", *Journal of Experimental Zoology (Molecular and Developmental Evolution)* 304B: 18-27.
- Simon, Herbert A. (1962), "The Architecture of Complexity", *Proceedings of the American Philosophical Society* 106: 467-482.

- Skipper, Robert A., and Roberta L. Millstein (2005), “Thinking About Evolutionary Mechanisms: Natural Selection”, *Studies in the History and Philosophy of Biological and Biomedical Sciences* 36: 327-347.
- Von Dassow, George, and Ed Munro (1999), “Modularity in Animal Development and Evolution: Elements of a Conceptual Framework for EvoDevo”, *Journal of Experimental Zoology (Molecular and Developmental Evolution)* 285: 307-325.
- Von Dassow, George, Eli Meir, Ed Munro, and Garrett M. Odell (2000), “The Segment Polarity Network is a Robust Developmental Module”, *Nature* 406: 188-192.
- Waters, C. Kenneth (1994), “Genes made molecular”, *Philosophy of Science* 61: 163-185.