

Abnormal light signals and the underdetermination of theory by evidence in astrophysics

Felipe A. Asenjo*, Sergio A. Hojman†, Niels Linnemann‡ and James Read§

Abstract

We investigate the propagation of certain non-plane wave solutions to Maxwell’s equations in both flat and curved spacetimes. We find that the effective signal velocity associated to such solutions need not be c and that the signal need not propagate along null geodesics; indeed, more than this, we find that the information encoded in the signals associated with such solutions can be substantially non-local. Having established these results, we then turn to their conceptual-philosophical-foundational significance—which, in brief, we take to be the following: (i) one should not *assume* that all electromagnetic waves generated in the cosmos are localised plane wave packages; thus, (ii) one cannot *assume* that signals reaching us from the cosmos arrive with a particular velocity (namely, c), and that such signals encode local information regarding their sources; therefore (iii) astrophysicists and cosmologists should be wary about making such assumptions in their inferences from obtained data—for to do so may lead to incorrect inferences regarding the nature of our universe.

Keywords: underdetermination, theory-dependence, electromagnetism, signal velocity, Airy beam

Contents

1	Introduction	1
2	Notions of signal propagation	2
2.1	Background on wave velocities	3
2.2	Velocities beyond isotropic waves	4
3	Warm-up: the Slepian wave	5
4	Maxwell equations in flat spacetime and FLRW cosmology	6
4.1	Maxwell equations in flat and curved spacetimes	6
4.2	Amplitude gradient effects	7
4.3	Polarisation gradient effects	9
5	The case of Airy beams	10
6	Foundational consequences	11
6.1	Underdetermination of theory by data	12
6.2	Theory-dependence of observation	13

1 Introduction

Einstein’s 1905 theory of special relativity, as presented in [Ein05], involves several key inputs: the relativity principle (that the laws of physics are the same in all coordinate systems in uniform translatory motion with respect to one another), the light postulate (that the speed of light is independent

*felipe.asenjo@uai.cl

†sergio.hojman@uai.cl

‡niels.linnemann@unige.ch

§james.read@philosophy.ox.ac.uk

of the speed of the source), the ‘standard’ Einstein-Poincaré clock synchrony convention, and the homogeneity and isotropy of space. From the relativity principle and light postulate, one can derive straightforwardly that the speed of light takes the same fixed value c in all inertial coordinate systems. So, in this sense, the constancy of the speed of light is close to being an input *assumption* of the special theory *circa* 1905—and, indeed, this constancy is often taken to be one of the defining features of the theory of relativity *tout court*—both special and general.

Note, however, that the lesson of Einstein’s 1905 paper was supposed to be the *kinematical constraint* that all laws of physics are conditioned so as to be invariant under Lorentz transformations. Indeed, Einstein would in later life lament the special significance afforded to light and electromagnetism in his 1905 presentation of the theory, opting instead to cut directly to this conclusion:

The content of the restricted relativity theory can accordingly be summarised in one sentence: all natural laws must be so conditioned that they are covariant with respect to Lorentz transformations. [Ein54]

However, even if one accepts (with the later Einstein) the above kinematical constraint as the ‘essence’ of special relativity (which also obtains locally in general relativity), one should avoid inferring too much from this constraint *vis-à-vis* the nature of propagating relativistic fields. For example: although it is true that plane wave solutions to the Maxwell equations do indeed propagate with determinate velocity c , it is in fact far from obvious that this continues to be the case (a) when one considers other solutions to said equations, or (b) one accommodates for the possibility of spacetime curvature. Indeed, for over half a century relativistic causality has in fact been acknowledged to be a delicate business: see [But07] for a recent survey of this terrain, and [AH17, LR21] for particular recent investigations into the behaviour of wavelike solutions to the Maxwell equations in curved spacetimes.

For these reasons, it is arguably more foundationally careful to regard Einstein’s 1905 theory of special relativity as taking the form of a *self-consistent bootstrap* (cf. [Cao01]): one begins with certain assumptions regarding the velocity of light; however, the conclusion at which one arrives (namely, the Lorentz invariance of the laws of physics) may—at least under certain circumstances—lead one to revise those very principles from which one began.

Our goal in the present article is to take up these issues, exploring in greater depth the propagation of certain non-plane wave solutions to Maxwell’s equations (in particular e.g. Airy wave solutions) in both flat and conformally flat curved spacetimes. We find that the effective velocity of such signals (*nota bene*: we will clarify different salient senses of wave velocity in what follows) need not be c and that such solutions need not propagate along null geodesics; indeed, more than this, we find that the information encoded in the signals associated with such solutions can be substantially non-local. Having established these results, we then turn to their conceptual-philosophical-foundational significance—which, in brief, we take to be the following: (i) one should not *assume* that all electromagnetic waves generated in the cosmos are localised plane wave packages (for why, indeed, should this be the case?); thus, (ii) one cannot *assume* that signals reaching us from the cosmos arrive with a particular velocity (namely, c), that they are effectively travelling on null geodesics, and that they encode local information regarding their sources; therefore (iii) astrophysicists and cosmologists should be wary about making such assumptions in their inferences from obtained data—for to do so may lead to incorrect inferences regarding the nature of our universe.

The structure of this article is as follows. In §2, we present the relevant background on waves and their velocities. In §3, we present a historical warm-up case regarding the unexpected behaviour of non-plane wave solutions to the Maxwell equations—namely, the case of the so-called Slepian wave. In §4, we consider more generally non-plane wave solutions to the Maxwell equations in both Minkowski and cosmological FLRW spacetimes, taxonomising different ways in which non-null or non-local propagation might arise. In §5, we focus in particular upon Airy solutions to the Maxwell equations in Minkowski and FLRW, and explore the propagation behaviour of these solutions. In §6, we discuss the philosophical and conceptual upshots of these results *vis-à-vis*, *inter alia*, empirical underdetermination of light signals received in astronomy and cosmology and the theory-dependence of observation.

2 Notions of signal propagation

In this section, we present the relevant background on (a) different notions of wave velocity (§2.1), and (b) velocities of non-plane waves (§2.2).

2.1 Background on wave velocities

Is it possible that electromagnetic waves *in vacuo* might ‘signal’ at some speed other than c ? *Prima facie*, there seem to be three distinct possible answers to this question:

- (i) It is impossible that electromagnetic waves could signal at some speed other than c .
- (ii) Electromagnetic waves’ signalling at some speed other than c is possible in principle, but not in practice. This would raise a conundrum—possibly of deeper significance—as to why exactly we do not see such signals.¹
- (iii) Electromagnetic waves’ signalling at some speed other than c is possible in practice. This, of course, would call for more research in this direction.

Before we can make progress in deciding which of the above three answers is correct, however, we need first to get a better handle on what is meant by ‘signal’, and what is meant by ‘signal speed’, in the context of wave propagation. A modest working definition for ‘signal’ is this: a propagating disturbance that encodes information where, standardly, the disturbance is taken to propagate locally. (The local nature of the propagation is an often tacitly assumed aspect which will become relevant later on.)

Disturbances can be modelled through wave packages. There are different ways to understand this:

- (a) The wave package *per se* is taken to represent the disturbance.
- (b) Parts of the wave packages are taken to represent a notion of disturbance (say, the high-frequency components).

Given this ambiguity, there are various distinct notions of signal speed which can be associated to a wave package, for example:

- (I) The group velocity, which is the velocity with which the overall envelope of the wave’s amplitudes propagates through space. (Regarding group velocity as the relevant notion of signal velocity would thereby be in line with (a) above.)
- (II) The front velocity, which is the velocity of the high frequency components of the wave (see e.g. [Sho03]). (Regarding the front velocity as the relevant notion of signal velocity would thereby be in line with (b) above.)

Now, it is general consensus that (1) the concepts of phase velocity or group velocity are generally not valid concepts of signal velocity (although, of course, they can in cases serve as proxies for it, especially if they all agree), and that (2) the concept of front velocity—the high-frequency limit of the phase velocities of a wave package’s components—serves as an upper bound to signal speed of a wave package, which, however, need not always be saturated (causal influences in water, for instance, propagate at speeds less than their front velocity): see [GMCB98, SGN03]. Beyond this, however, the notion of the (signal) velocity of a wave is a delicate matter, and must be assessed on a case-by-case basis.

One other notion of wave velocity which will be of particular relevance in the context of our investigations in this article is the transverse ‘velocity’ vector $\vec{v} = \vec{\nabla}\phi/\omega$, where ϕ is the phase of the wave and ω its frequency (this will be explained in more detail in §4.3)—such a notion of wave velocity is used in ongoing optics research [NEC⁺18, NNB22, PN98], and has the merit of being associated directly to the normalised Poynting vector of an electromagnetic wave (and thus clearly of significance to energy—and in this sense signal—propagation via the wave). We will, indeed, have several occasions to compute this quantity in later sections of this article.

¹This second option would certainly appear to be in conflict with Gell-Mann’s ‘totalitarian principle’: *everything not forbidden is compulsory* [GM56]. The issue is, perhaps, of a similar kind to that of why we see only retarded but not advanced solutions in electrodynamics.




Figure 1: Wave fronts for a plane wave.




Figure 2: Discontinuous wave fronts for a generalised ‘wave’.

2.2 Velocities beyond isotropic waves

Standardly, the aforementioned wave velocities are defined only with respect to plane waves or spherical waves. This, however, commits one to a basic ontology of waves whose fronts fulfil certain standards of isotropy; one might wonder as to the extent to which such notions of wave velocities can be extended beyond the plane or spherical wave in particular. To make progress in answering this question, recall first that a plane wave $\psi(\vec{x}, t)$ can be written in the form

$$\psi(\vec{x}, t) = A(\vec{k}) \exp(\vec{k} \cdot \vec{x} - \omega t), \quad (1)$$

where \vec{k} is a constant three-vector known as the ‘wave number’, and ω is again constant, and denotes the frequency of the wave; moreover, the amplitude A of the wave is constant.

One might, however, seek to generalise *beyond* the notion of a plane wave. For example, consider generalised waves of the form

$$\chi(\vec{x}, t) = A(\vec{k}, \omega) \exp(\vec{k}(\vec{x}, t) \cdot \vec{x} - \omega(\vec{x}, t)t); \quad (2)$$

in this case, the wave number and frequency are spacetime-dependent quantities; the amplitude—which is a function of wave vector and frequency—is implicitly dependent on spacetime quantities through its dependence upon $\vec{k}(\vec{x}, t)$ and $\omega(\vec{x}, t)$. One might object that such a liberalisation of the notions of wave vector and frequency goes against the spirit of introducing the wave vector and frequency to begin with; after all, on a plane wave approach, wave vector and frequency parameterise a specific plane wave and thereby allow for the packaging of specific plane waves in a weighted manner through a wave vector- and frequency-dependent amplitude. However, even if each wave vector is spacetime-dependent, it is not the case that different waves cannot be summed; in this sense, the generalisation (2) is perfectly legitimate. Needless to say, the generalised waves could be decomposed further into plane waves proper. But the point of the general wave form is to allow for the setting up of notions of wave velocity that are not limited to plane waves *specifically*, independent of the mathematical fact that (as is of course well-known) plane waves form a basis for the solution space for a wave equation.




Figure 3: Continuous but spacetime-dependent wave fronts for a generalised wave.

Physically, the wave number picks out level surfaces in space which correspond to wave crests: for the canonical plane wave, these are evenly-spaced, as per Figure 1. In the more general case in which the wave number is an arbitrary function of spacetime, however, it may manifest discontinuities and caustics, as in Figure 2. In that case, it is by no means clear that notions such as phase, group, or front velocity will be well-defined mathematically. Therefore, if one is to continue to make sense of such notions of wave velocities once one has generalised beyond the plane wave in the above manner, it is prudent to impose minimal restrictions upon the wave number—in particular, it is prudent to impose that its level surfaces be *continuous* (or maybe even smooth) in space. In this way, the wave vector can nevertheless pick out ‘non-plane wave’ surfaces, such as those of Figure 3. In a like manner, in order for notions of phase, group, and front velocity to be well-defined in the general case, one of course will also require that the wave frequency ω be continuous, for essentially the same reasons as above.

With these points in mind, we now have the resources in order to assess the salient velocities of solutions to the Maxwell equations beyond plane waves: it is to these issues that we now turn.

3 Warm-up: the Slepian wave

As a warm-up to our discussion in later sections of the propagation behaviour of various solutions to the Maxwell equations in certain spacetimes (specifically, Minkowski and FLRW), let us consider first in this section a somewhat infamous proposal for faster-than-light signalling from electromagnetic waves made by Slepian [Sle49].² Slepian gave the following solution to the free space wave equation for the electromagnetic vector potential \mathbf{A} in flat spacetime, $\nabla^2 \mathbf{A} = \frac{1}{c^2} \frac{\partial^2 \mathbf{A}}{\partial t^2}$, in Coulomb gauge $\nabla \cdot \mathbf{A} = 0$:

$$\mathbf{A} = \cos[k(x - vt)] \cos(k\Gamma y) \hat{\mathbf{z}}, \quad (3)$$

where $\Gamma := \sqrt{\frac{v^2}{c^2} - 1}$. The surprising feature of the solution (3) is that the waveform seems to move in x -direction at a speed $v > c$ (required in order for Γ to be real)—i.e., faster than the speed of light.

McDonald [McD18] has provided a sober analysis of Slepian’s “academic puzzler” by decomposing the vector potential into two plane waves, as

$$\mathbf{A} = \frac{A}{2} \exp i(k_+ r - \omega t) \hat{\mathbf{z}} + \frac{A}{2} \exp i(k_- r - \omega t) \hat{\mathbf{z}}, \quad (4)$$

where $\omega = vk$ and $\mathbf{k}_\pm = k\hat{\mathbf{x}} \pm k\Gamma\hat{\mathbf{y}}$. Clearly, the phase velocity for each of these waves is c :

$$v_\pm = \frac{\omega}{|\mathbf{k}_\pm|} = \frac{\omega}{k\sqrt{1 + \Gamma^2}} = \frac{vk}{k(v/c)} = c; \quad (5)$$

however, their directions are different. Thus, the Slepian wave can be understood as the result of two plane waves emitted from distinct sources—in this sense, it encodes *non-local* information (we return to this issue in the following sections). In any case, as McDonald goes on to point out, the Slepian

²In fact, the extent to which Slepian himself wanted his proposal to be read as one for arbitrary signalling speeds is ambiguous—his article ends with the passage: “We conclude then that electromagnetic waves in free space exist which travel respectively with any and all velocities, v . How about this?” Nevertheless, we cannot help but think that he *must* have meant a substantial sense of ‘travelling’, tantamount to signal propagation.

wave is a completely legitimate solution to Maxwell’s equation (in fact, it also goes under the name of the TE_{10} mode of rectangular waveguides)—but it simply cannot be used for faster-than-light *signal* propagation any more than the travelling intersection point of a scissors’ blade: a signal that is sharp in frequency—such as the Slepian wave—has to be modulated across the *whole wavefront*. Strictly, however, this takes infinite time. Thus, the combination of wavefront encoding and actual propagation of the wavefront of a Slepian wave prohibits faster-than- c propagation.³ Note also, finally, that, since they are plane waves, the phase and group velocities of the constituents of the Slepian wave in (4) will also be c .

4 Maxwell equations in flat spacetime and FLRW cosmology

In this section, we present two solutions for abnormal light propagation arising from the vacuum Maxwell equations in flat spacetime and in Friedman-Lemaître-Robertson-Walker (FLRW) cosmology. In particular, we show that this abnormal light propagation involves beam constituents which move on bent rather than straight trajectories—and so not on null geodesics. These can occur for two different reasons: (i) the existence of second-order amplitude wave gradients (giving rise to the known ‘Bohm potential effects’), and (ii) the existence of second-order polarization wave gradients. Both are discussed below, with corresponding particular solutions.

The structure of the section is this. In §4.1, we remind the reader of the relevant details of the Maxwell equations in curved spacetimes. In §4.2, we consider amplitude gradient effects. In §4.3, we consider polarisation gradient effects.

4.1 Maxwell equations in flat and curved spacetimes

In general, the Maxwell equations in vacuum flat spacetime for the electric field \mathbf{E} and the magnetic field \mathbf{B} , are written (in vectorial notation) as

$$\begin{aligned}\nabla \cdot \mathbf{B} &= 0, & \nabla \cdot \mathbf{E} &= 0, \\ \partial_t \mathbf{E} &= \nabla \times \mathbf{B}, & \partial_t \mathbf{B} &= -\nabla \times \mathbf{E}.\end{aligned}\tag{6}$$

The abnormal propagation of light in this context (i.e., *even in flat spacetime*) has been studied theoretically and experimentally as solutions of these equations for electromagnetic waves (see *inter alia* [SBDC07, SBDC08, BNKS14, KBNS12, PBBS18, Hac11, AMCK⁺16, CE13, SC07, CRNW10, PCP⁺16, ZBH13]).

Before we consider this in more detail, however, let us first demonstrate that in FLRW cosmologies, electromagnetic wave propagation can in fact be described by a similar set of Maxwell equations. As is well-known, the Maxwell equations in curved spacetime can be written

$$\nabla_\mu F^{\mu\nu} = 0, \quad \nabla_\mu F^{*\mu\nu} = 0,\tag{7}$$

written in terms of compatible covariant derivative operators for a general spacetime metric $g_{\mu\nu}$. As usual, we can now define the vectorial electric and magnetic fields as $E_i = F_{i0}$, $D^i = \sqrt{-g}F^{0i}$, $B^i = \varepsilon^{0ijk}F_{jk}$ and $\varepsilon^{0ijk}H_k = \sqrt{-g}F^{ij}$, with relations

$$\begin{aligned}D^i &= -\sqrt{-g}\frac{g^{ij}}{g^{00}}E_j + \varepsilon^{0ijk}\frac{g_{0j}}{g_{00}}H_k, \\ B^i &= -\sqrt{-g}\frac{g^{ij}}{g^{00}}H_j - \varepsilon^{0ijk}\frac{g_{0j}}{g_{00}}E_k.\end{aligned}\tag{8}$$

Then, the Maxwell equations can be written

$$\begin{aligned}\nabla \cdot \mathbf{B} &= 0, & \nabla \cdot \mathbf{D} &= 0, \\ \partial_0 \mathbf{D} &= \nabla \times \mathbf{H}, & \partial_0 \mathbf{B} &= -\nabla \times \mathbf{E}.\end{aligned}\tag{9}$$

The above system is general for any spacetime. However, in what follows let us now consider specifically the FLRW metric

$$g_{\mu\nu} = (-1, a^2, a^2, a^2),\tag{10}$$

³This issue has been familiar since at least the work of Brillouin [Bri60].

where $a = a(t)$ is the usual FLRW scale factor. In this case, relations (8) reduce to

$$\mathbf{D} = a\mathbf{E}, \quad \mathbf{B} = a\mathbf{H}, \quad (11)$$

and then, the above Maxwell equations (9) become

$$\begin{aligned} \nabla \cdot \mathbf{B} &= 0, \quad \nabla \cdot \mathbf{D} = 0, \\ \partial_\tau \mathbf{D} &= \nabla \times \mathbf{B}, \quad \partial_\tau \mathbf{B} = -\nabla \times \mathbf{D}, \end{aligned} \quad (12)$$

—i.e., just like the free Maxwell equations in flat spacetime!—by defining the conformal time

$$\tau = \int \frac{dt}{a}. \quad (13)$$

Therefore, everything that is fulfilled for electromagnetic waves in free flat space occurs also for cosmological FLRW electromagnetic waves, in the conformal time τ . In flat spacetime, $a = 1$ and $\tau = t$. In a sense, of course, this is unsurprising, since (a) the FLRW solution is conformally flat, and (b) the source-free Maxwell equations are conformally invariant, so any solution of said equations in Minkowski spacetime will remain a solution in FLRW spacetime. In this sense, the experimental results that demonstrate that abnormal (i.e., non-geodesic) light propagation is possible in flat spacetimes is unavoidable imply also that the same kinds of propagation arise in cosmological solutions.

With the set of equations (12) in hand, we find that the magnetic field \mathbf{B} follows the general wave equation

$$(\partial_\tau^2 - \nabla^2) \mathbf{B} = 0. \quad (14)$$

(A similar equation can be found for the electric field \mathbf{E} ; in the following we focus for simplicity only on the wave equation for \mathbf{B} .) Usually, (14) is solved in terms of plane waves travelling to the speed of light (or, in relativistic vocabulary, on null geodesics). However, this wave equation has other possible solutions that present non-trivial behaviour. As an exemplification of this, in the following two subsections we consider two of these solutions—in the first case, the solutions involve second-order amplitude gradients; in the second, they involve second-order polarisation wave gradients. Note, to repeat, that both of these classes of solutions (for the reasons articulated above) are solutions both of the Maxwell equations in flat Minkowski spacetime *and* in the (curved) FLRW spacetime which we take to represent accurately the large-scale structure of our universe.

4.2 Amplitude gradient effects

When the wavepacket solution for electromagnetic wave propagation is allowed to have amplitude gradients, then it modifies its propagation. This has been reported extensively in the context of the paraxial and non-paraxial propagation of light in both flat and curved spacetime geometries [BNKS14, PBBS18, KBNS12]. In these spacetimes, it has been shown that such wavepackets can follow non-geodesic trajectories, and indeed follow curved trajectories due to the interaction of amplitude properties of the wave with its medium.

In this subsection, we present the simplest case of such effects for a cosmological scenario; in particular, we shown that an electromagnetic wave can propagate in an accelerated fashion due to amplitude gradient effects. We look for a solution in the form $\mathbf{B} = B(u, y) \exp(iv)\hat{z}$, with only one component (with $u = x - \tau$ and $v = x + \tau$). In this case, the wave equation (14) becomes

$$(-4i\partial_u + \partial_y^2) B = 0. \quad (15)$$

This is a Schrödinger-like equation, which has several known solutions. One of them is in the form of an Airy function [BB79, Lek09]. Therefore, in our case, the field can propagate as

$$B(\tau, x, y) = B(u, y) = \text{Ai} \left(2y - u^2 - iau \right) \exp \left(ay - au^2 - i\frac{a^2}{4}u - 2iuy + \frac{2}{3}iu^3 \right), \quad (16)$$

where Ai is the Airy function, and a is an arbitrary parameter that allow us to have finite energy of the wavepacket [Lek09].

The above solution presents curved trajectories in the y - u plane, i.e., as the wavepackets propagate along $u = x - \tau$ direction, it deflects also on transversal y -direction. This behaviour does not occur




Figure 4: Contour plot, in the y - u plane with $a = 0.1$, of the square of intensity of wavepacket (17) for the different lobes of the solution. The red line shows the deflection of this maximum intensity of this solution on this space.

for a plane wave, which does not manifest any kind of deflection in its transverse direction. This can be studied numerically through the intensity of this wavepacket

$$b = \sqrt{\mathbf{B}^* \mathbf{B}} = \sqrt{B^* B}. \quad (17)$$

In Fig. 4, we contour plot the square of intensity b^2 , on the y - u plane with $a = 0.1$, to show that this Airy solution has several lobes that behave differently, deflecting in a different manner. In the plot, the red line shows the curved trajectory of the maximum intensity of this wavepacket. In the usual plane wave solution, all trajectories should be vertical lines in such plots. Therefore, the abnormally-propagating solution (16) deflects itself during its propagation due to its dynamics on the transverse plane.

This can also be studied via the Poynting vector of this wave. This vector establishes the direction of the energy flux of the wave. The Poynting vector is $\mathbf{S} = \frac{1}{2}(\mathbf{D}^* \times \mathbf{B} + \mathbf{D} \times \mathbf{B}^*)$, where \mathbf{B} is the already-calculated magnetic field, and $\mathbf{D} = \int d\tau \nabla \times \mathbf{B}$. In the typical case of plane waves (with constant amplitudes), the only non-zero component of the vector field \mathbf{D} is in the y -direction, and the flux of energy (Poynting vector) is in the x -direction. However, in this case,

$$\mathbf{D} = D_x \hat{x} + D_y \hat{y} = \int d\tau \partial_y (B e^{iv}) \hat{x} - \int d\tau \partial_x (B e^{iv}) \hat{y}, \quad (18)$$

and the Poynting vector has a component in the y -direction as well, $\mathbf{S} = S_x \hat{x} + S_y \hat{y}$, with

$$\begin{aligned} S_x &= \frac{1}{2} (D_y^* B e^{iv} + D_y B^* e^{-iv}), \\ S_y &= -\frac{1}{2} (D_x^* B e^{iv} + D_x B^* e^{-iv}). \end{aligned} \quad (19)$$

This is due only to the transversal (extended) spatial dependence of the amplitude of the wave, which now depends also on the y -direction. This implies that for this wave solution the energy flux follows a curved trajectory in the y - u plane.

4.3 Polarisation gradient effects

To compliment the above-discussed situations in which the propagation of electromagnetic waves can be affected by an amplitude gradient, we ask in this subsection whether effects on *polarisation* can likewise induce abnormal wave propagation. In general, the wave equation (14) describes the behaviour of polarised waves, and therefore more general solutions than the plane wave can again be identified. For example, recently it has been shown (both theoretically and experimentally) in [NNB22] that second-order gradients on the phase of polarised waves in vacuum flat spacetimes can also induce beam deflections on the wave. Thus, we expect that similar effects occur for cosmological light.

Consider a magnetic field with a transverse polarized part in the form

$$\mathbf{B}(\tau, x, y, z) = \rho^{1/2} [\cos \gamma \hat{x} + \sin \gamma \hat{y}] \exp(i\phi + i\omega\tau) + B_z \hat{z}, \quad (20)$$

where ρ is the amplitude of the wave, γ is the polarization angle, ϕ is total wave phase, and ω is the wave frequency. The physical quantities ρ , γ , and ϕ can have full dependence on (conformal) time and space coordinates. The frequency ω is a constant for this solution. B_z denotes a longitudinal part of the field. Using this ansatz in (14) for the transverse part, we obtain the four equations

$$\begin{aligned} \nabla \cdot (\rho \nabla \gamma) &= 0, \\ \nabla \gamma \cdot \nabla \phi &= 0, \\ \nabla \cdot (\rho \nabla \phi) &= 0, \\ |\nabla \phi|^2 - \omega^2 - \frac{\nabla^2 \sqrt{\rho}}{\sqrt{\rho}} + |\nabla \gamma|^2 &= 0. \end{aligned} \quad (21)$$

On the other hand, B_z must fulfil (14). In principle, we see already from this (in particular from the last equation) that the gradient of the polarization angle can induce a deflection on the trajectory of the wave, calculated through the second order gradient of the phase $\nabla(|\nabla \phi|^2)$ [NNB22].

A particularly neat example of this effect can be obtained for a perturbative solution. Let us suppose that

$$\begin{aligned} \rho(z) &\approx \rho_0 + 2\epsilon \rho_1(z), \\ \gamma(x, y) &\approx \epsilon^{1/2} \gamma_1(x, y), \\ \phi(x, y, z) &\approx \omega z + \epsilon \phi_1(x, y, z), \end{aligned} \quad (22)$$

where ϵ is the small parameter that we use to keep track of the perturbative order in the equation. Here ρ_1 , γ_1 , and ϕ_1 are the perturbations, while ρ_0 is a constant. Thereby, the transverse part of this solution is a polarised perturbation of a plane wave travelling in the z -direction.

The first and second equations of set (21) (both at $\mathcal{O}(\epsilon^{1/2})$) allow us to establish that

$$\gamma_1(x, y) = \omega \alpha x y, \quad (23)$$

where α is a constant. In addition, the third and fourth equations (both at $\mathcal{O}(\epsilon)$) allow us to find that

$$\begin{aligned} \rho_1(z) &= \frac{\alpha^2}{2} z^2, \\ \phi_1(x, y, z) &= \frac{\alpha^2}{2\omega} z (1 - \omega^2(x^2 + y^2)). \end{aligned} \quad (24)$$

Finally, from constraint (12), the above transverse solution for the ansatz allows us to find B_z at order $\mathcal{O}(\epsilon^{1/2})$. This is

$$B_z(\tau, x, z) = i\epsilon^{1/2} \sqrt{\rho_0} \alpha x \exp(i\omega z + i\omega\tau), \quad (25)$$

which also solves (14) identically.⁴

From the above solution we can see that small changes on the polarization angle will induce small changes on the wave trajectory. The wave trajectory velocity \mathbf{v} can be obtained through the gradient

⁴We are not aware of an example which fully isolates the polarisation gradient effect from the amplitude gradient effect.

of its phase $\mathbf{v} = v_x \hat{x} + v_y \hat{y} + v_z \hat{z} = \nabla \phi / \omega$ [NNB22] (on this notion of velocity, recall our discussion in §2.1). Thus, for example, this wave packet satisfies

$$\left| \frac{dv_y}{dz} \right| = \epsilon \alpha^2 y, \quad (26)$$

implying that, as the wave propagates longitudinally in z -direction, it deflects along the transverse y -direction due to the polarization gradient effect. Similar behaviour happens for the x -direction.

As before, this can be studied using the Poynting vector of this solution. Its complete calculation is highly involved and not very illuminating, but instead one can obtain the different ϵ order corrections for the solution. For the case in hand, the Poynting vector values are $\mathbf{S} = \frac{1}{2}(\mathbf{D}^* \times \mathbf{B} + \mathbf{D} \times \mathbf{B}^*) = S_x \hat{x} + S_y \hat{y} + S_z \hat{z}$, where

$$\begin{aligned} S_x &= \mathcal{O}(\epsilon^{3/2}), \\ S_y &= \epsilon \alpha^2 y z \rho_0 + \mathcal{O}(\epsilon^{3/2}) \\ S_z &= -\rho_0 + \epsilon^{1/2} \frac{\alpha \rho_0}{\omega} + \epsilon \frac{\alpha^2}{2} (\rho_0(x^2 + y^2) - 2z^2 - \rho_0/\omega^2) + \mathcal{O}(\epsilon^{3/2}), \end{aligned} \quad (27)$$

implying that the main flux of energy occurs in the z -direction (which is the expected result as it is the non-perturbative plane wave solution), along with several corrections. However, we find that, at ϵ order, energy also flows in the y -direction. Thus, while the wave mainly propagates in z -direction, it also deflects in y -direction. This is in agreement with the velocity gradient (26), implying that $\mathbf{v} = \nabla \phi / \omega$ plays the role of the normalised Poynting vector of this electromagnetic wave.

5 The case of Airy beams

So far, we have seen that both amplitude and polarisation gradient effects can affect the propagation trajectories of electromagnetic waves in both flat and FLRW spacetimes. In this section, we explore the physical status of optical Airy beams, the discussion of which will also help us to understand better the status of the specific Airy-type solutions in curved spacetimes discussed the previous section. Generally from now on, we will use Airy beam scenarios as paradigmatic cases for abnormal light propagation (as by now familiar from wave propagation in dielectric media), but this focus is in no way intended to suggest that abnormal propagation is limited only to Airy beam behaviour.

Airy beams are usually associated to solutions of the paraxial approximation of the Helmholtz equation. That is, the Helmholtz equation $\nabla^2 f + k f = 0$ with ansatz $A(r) = u(r) \exp(ikz)$ is approximable to

$$\nabla_{\perp}^2 u + 2ik \frac{\partial u}{\partial z} = 0 \quad (28)$$

provided that $|\frac{\partial^2 u}{\partial z^2}| \ll |k \frac{\partial u}{\partial z}|$.⁵ Apparent core features of Airy beams include that they are weakly diffracting (amplitude profile stays close to invariant), curve in propagation (‘parabolic shape’) and can be self-healing (the profile can recover its original form when obstacles are placed in the path of their main lobe). All these three aspects of behaviour can be explained by way of a detailed analysis of Airy beams in terms of a ray decomposition approach: see [KH10]. Interestingly, Airy beam solutions to the paraxial approximation of the Helmholtz equation are the only diffraction-free solutions in one transverse-dimension. This, however, leaves open the possibility for non-diffraction-free paraxial accelerating waves or those with more than one transverse direction.

What about the non-paraxial case? Consider the 2d case:

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial z^2} + k^2 f = 0, \quad (29)$$

As reviewed in [ECSC19], so-called half-Bessel beams are diffraction-free solutions to the 2d Helmholtz equations. Notably, they can follow circular, ellipsoidal and other paths, whereas the Airy beams—solutions to the paraxial approximation of the 2d Helmholtz equation—trace out parabolic paths.

⁵Notably, the equation is mathematically equivalent to the Schrödinger equation—thus the appearance of Airy waves in quantum mechanics too (albeit there they are ‘over time and space’ and not just space).

What then is the significance of Airy beams *qua* physical signals? The work of [KH10], which analyses Airy beams in terms of geometric rays, is especially helpful in answering this question. The set-up is as follows: the non-paraxial equation is considered in two dimensions,

$$\partial_x^2 u + 2ik\partial_z u = 0, \quad (30)$$

together with an initial field distribution in the aperture plane $z = 0$ of form $u(z = 0) = \text{Ai}(x/x_0)$. The radiated field $f = u \exp(ikz)$ is then given by Kirchhoff-Huygens integral

$$U(r) = 2 \int_{-\infty}^{\infty} dx' \phi_0(x') \partial_{n'} G(r, r') \quad (31)$$

with Rayleigh-Sommerfeld diffraction Green function G . One can distinguish between contributions to the field U from the main lobe of the aperture (call this U_1) and contributions from the remainder of the aperture (call this U_2). The analysis of [KH10] then leads to the *prima facie* surprising insight that the main curved lobe of the Airy beam “is not generated by the first main lobe of the aperture field, but by some other part” (p. 2). In particular, it is concluded in light of the analysis that an Airy beam is *not* a beam in the sense of “a local wave function whose evolution along its propagation trajectory is described by a local self-dynamics” (p. 2).

However, the results of [KH10] need to be put into context: it is true that Airy beams exhibit a straight line movement of their intensity centroid, but the shape of their intensity structure follows a curved path. It is the field profile of the beam, rather than the center of mass, that has immediate influence on optical processes. As for instance stressed in the review [ECSC19], “the majority of optical processes like nonlinear effects, *detection*, imaging etc., are directly dictated by the field profile itself, not by the ‘center of mass’ of the beam, which is a mean property (averaged over the beam cross section).” (p. 686, our emphasis).⁶ Generally speaking, as [PBBS18] discuss, the main feature of accelerating beams carrying finite power is that their main lobe follows a curved path while their center of mass does follow a straight trajectory. Again, this has significant consequences for light-matter interactions, as all such interactions depend on the local intensity of the beam rather than its center of mass; in fact, this feature has enabled a variety of applications, including curved plasma channels, manipulation of microparticles, laser micromachining, single-molecule imaging, and light-sheet microscopy, among others.

6 Foundational consequences

Up to this point, we have made the case that once one generalises beyond plane wave solutions to the Maxwell equations in both flat and curved spacetimes, the propagation of the waves may be affected so as to lead to non-null motion; moreover, the case can be made that some of these solutions encode non-local information (recall again for instance the Slepian wave).

Prima facie, this has the potential to be a profound result, given that the standard assumption in practising astrophysics and cosmology is that electromagnetic signals received at Earth travel on null geodesics, and are localised entities. In particular, given the cornucopia of different ways in which wave-like solutions to the Maxwell equations have been generated in the lab as well as brought into application, the assumption that *all* propagating solutions to the Maxwell equations in the cosmos resemble plane wave solutions begins to look not only miraculous, but also implausible.⁷ Airy beams in the sense of §4.2 can for instance be implemented straightforwardly given that the Fourier transform of the Airy wave is a Gaussian beam with an additional cubic phase—for instance by using a spatial light modulator [ECSC19, p. 687]⁸ or specially-designed lenses [YSA10, PSAT10] on a Gaussian laser mode; polarised beams in the sense of §4.3 can in principle be generated using an initial polarized light beam that passes through spatial light modulators and polarization filters [NNB22].⁹

⁶On p. 687, the authors also write: “The study of optical accelerating waves effectively began with the understanding that what really matters is not the trajectory of the ‘center of mass,’ but the curved evolution of the field itself. Thus, a truncated Airy beam (which is physically realizable since the beam must have finite power), interacts with its surroundings—whether they are particles or other waves—according to its accelerating field structure, not according to its centroid, which moves instead on a straight line so as to ensure the preservation of transverse momentum.”

⁷Recall the response to option (ii) in §2.1, discussed in footnote 1.

⁸See also [SBDC08, BSDC08].

⁹We thank Zhigang Chen and Nikos Efremidis for correspondence on these issues.

Again, one would thus arguably expect likewise that both types of abnormal light propagation could be generated through processes at the astrophysical scales. In response to this expectation, one might argue that such abnormal light propagation would only arise in cosmological scenarios as the result of finely-tuned set-ups which are unlikely to occur naturally. Our personal view is that as-of-yet there is insufficient evidence to warrant this armchair prohibition—but we refrain from a final verdict on these matters pending further empirical investigation; the foundational consequences explored in the following are to be regarded as being contingent on an affirmative answer to this existence question.

In the remainder of this section, we explore the philosophical-conceptual-foundational significance of the (assumed) existence of such non-standard electromagnetic wave solutions in cosmological scenarios with respect both to the underdetermination of theory by data (§6.1), and with respect to the theory-dependence of observation (§6.2).

6.1 Underdetermination of theory by data

Underdetermination of theory by data is often posed in the philosophy of science in a ‘global’ manner: if two different theories make the same empirical predictions, then how is one to tell which of those theory ‘truly’ describes reality? Issues of underdetermination, however, can of course also be a ‘local’ matter:¹⁰ given such-and-such empirical data/evidence, which theoretical hypotheses are compatible with that data/evidence?¹¹

Clearly, it is this latter notion of underdetermination of theory by data which is relevant to our points in this article. In particular, given the possibility of abnormal wave propagation in curved spacetimes as explored in the foregoing sections of this article, there arises a novel issue of underdetermination by data *vis-à-vis* the astronomical electromagnetic signals which we receive on (or around, via satellites) the Earth. In particular—and this in addition to the fact that the trajectories of photons are well-known to bend in light of the curvature of spacetime in general relativity—

1. Are such signals truly propagating on null geodesics and with determinate velocity c ?
2. Are such signals in fact encoding information about non-local sources (*à la* the Slepian wave)?

Given that such signals are in the relevant astrophysics and cosmology communities almost invariably *assumed* to propagate locally on null geodesics, there is—in light of the above—a pressing need to confront these questions.

In particular, in an astrophysical context, the roughly-detected beam segments as we register them on or around the Earth seem to leave open whether they are part of standard beam originating in sources ‘straight ahead’, or are better fitted to a model of abnormal wave propagation that is associated to a different local source (or maybe even to various sources at once).

This issue of underdetermination arising from the possibility of abnormal wave propagation is reminiscent of how gravitational lensing for light was traditionally regarded as a source for underdetermination by data of astrophysical phenomena. There is, for instance, the concrete suggestion by Barnothy and Barnothy from 1965 [BB68] that quasars are Seyfert galaxies subject to gravitational lensing rather than a class of cosmic objects of their own. Notably though, by now one can fairly well decide between whether or not gravitational lenses have influenced a beam’s path—the theory behind lensing has evolved and given rise to decisive observational traces of lensing in the beam [And15].

The currently-presented issue of underdetermination might be resolved in a similar manner at some point in the future. In this context, it is interesting to note the fair amount of optimism in fields of ‘historical’ research (to which astrophysics and cosmology can be counted to the extent that they make sense of past data) to the effect that apparent issues of underdetermination by data can eventually be resolved (and thus are in particular not issues of in-principle underdetermination). Consider, for example, the following passage from [And15] (echoing [Cle02]):¹²

¹⁰What we mean here is essentially tracking the difference between ‘strong’ versus ‘weak’ underdetermination, as summarised in e.g. [Lad01].

¹¹In what follows, we elide some philosophical subtleties by using ‘data’ and ‘evidence’ interchangeably.

¹²Cf. the optimism about the historical sciences defended by Currie [Cur18]. Here is Cleland on this issue:

Historical researchers investigating particular past events cannot test their hypotheses by performing controlled experiments. But this doesn’t mean that they cannot procure empirical evidence for them. Because of the asymmetry of over-determination, there are usually an enormous number of subcollections of the effects of a past event that are individually sufficient (given the right theoretical assumptions) to infer its occurrence. The trick is finding them. Many of these overdetermining traces (e.g., a splinter of glass, a

[I]f an event has occurred, it is very difficult to make things look like nothing had happened. The situation seems analogous to criminal cases, where it is reasonable to hope that the culprit has left some traces that will make it possible to identify him. In this sense, it is difficult to fake past events, i.e. to create all expectable traces artificially without the occurrence of the actual event. While in an experimental approach, the underdetermination problem makes it difficult to find the “true” (responsible) causes for an observed outcome, the richness of traces in historical science cases can help to distinguish true causes from faked causes. The large number of effects that an event usually creates leads to a so called “overdetermination of causes”. (p. 6)

So far, so good—but what about now? How strong really is the case of underdetermination raised by abnormal signal propagation? Are there genuine, concrete cases in which explicit alternative hypotheses for an arriving wave package in terms of non-geodesic (possibly even non-localised) sources are known or can be found?

In the remainder of this subsection we want to make progress on this complex of questions by comparing the situation of beam detection in astrophysics to that in a lab. The difference between beam detection in astrophysics and in a standard lab scenario is striking: while in the latter case the full extent of the beam can be studied from various perspectives (using all sorts of manipulations and measurement across the full length of the beam), in the former case the beam path is only known around its detection point (which is, given the distance the beam usually has travelled in an astrophysical context, a rather insignificant segment of the overall beam path). Therefore—much more than in the laboratory—in astrophysics theoretical background assumptions are required for making a statement on the full beam path. It is here that the results on abnormal light propagation in curved spacetime come in: they call into question the background assumption that wave signals follow the null geodesics of the background geometry; more than that: concrete alternatives for long-term beam propagation are provided (say in the form of specifically bent Airy wave packages), albeit only with respect to beam’s *path*.

Note though that these concrete alternative hypotheses for the beam paths are not *prima facie* part of a larger overall causal account behind the recorded beam data. In particular, it is not clear at this initial stage of hypothesising that the alternative beam path can really be due to a sensible physical source. Consider the case that the abnormally behaving beam is due to a highly non-localised source (as seems to be the case at least for the Airy beam solution on Kaganovsky’s analysis [KH10]). There are two options: (1) the alleged beam is effectively noise resulting from overlying locally sourced beams; then, it would, however, be a highly conspiratorial matter that the alleged beam came into being in any case—especially if we detect a whole class of beams of this form rather than just a single one. Or, (2) the beam is the result of some genuine physical mechanism which, however, has to be of non-local nature. Given that (1) and (2) are both rather unattractive options methodologically, it seems as if the standard hypothesis that (0) the detected beam moves on null geodesics (with a well-defined local source)—when available—is preferable.

In any case, and picking up the expectations mentioned above of an eventual resolution of the underdetermination issue, one should be able to decide by crucial experiment between specific abnormal Airy wave solutions and standard (geodesic) beam solutions: the characteristic mark of the abnormal Airy wave solution is a specific polarisation of light that should be subject to detection, even locally.¹³ Likewise, it is to be expected that other forms of abnormal light propagation will have signatures that can ultimately make them distinguishable from (say) Gaussian beams.

6.2 Theory-dependence of observation

Given the assumption that light propagates on null geodesics which is at least implicit (and, indeed, sometimes explicit) in most reasoning in astrophysics and cosmology, those physicists’ descriptions of

footprint) occupy small, local regions of space, bringing them within the limited range of human sensory experience. This places scientists investigating the remote past in the position of criminal investigators. Just as there are many different possibilities for catching a criminal, so there are many different possibilities for establishing what caused the demise of the dinosaurs, the origin of the universe, etc. Like criminal investigators, historical scientists collect evidence, consider different suspects, and follow up leads. (p. 490)

¹³On ‘crucial experiments’ more generally, see [Lak74].

observed results inevitably become a function of that assumption. For example, we hear regularly—on the basis of the assumption—that such-and-such galaxy is such-and-such many millions of light years away. Although such statements might appear to be matters of brute empirical fact, the above discussion should make clear that they are in fact much more theory-laden than they might initially appear.

Clearly, the central point of the present article is that we should be open to revising assumptions of this kind—if, on the basis of more detailed considerations (say, to do with how said electromagnetic waves are generated in such-and-such astronomical body, e.g. a star, or filtered in such-and-such nebula), we come to appreciate that we are not dealing with plane waves, and that (rather) the signals under consideration need not manifest geodesic motion, or need not encode only local effects, then in that case our descriptions of what we are observing on the basis of received signals will change. It seems to us correct to state that a process of reflective equilibrium is an appropriate way to characterise this process: (i) astrophysics (in this case) begins with certain general and univocal assumptions (e.g., to do with the null propagation of light), then (ii) uses those assumptions to gather data on astrophysical bodies (e.g. stars) and to develop a more sophisticated physics of the behaviour of those bodies, but (iii) that very understanding might lead us to revise the very assumptions from which we began (in this case, to do with the propagation of light emitted from such bodies). In turn, this only goes to illustrate the complex interrelations between the theoretical and the empirical in modern science; the lesson, of course, is that we should not (and need not) be too keen to hold onto established consensus, if theoretical considerations on the basis of amassed empirical evidence (which might have been gathered from within the framework of that very consensus) speak against it.¹⁴

Acknowledgements

We are grateful to Zhigang Chen and Nikos Efremidis for helpful discussions. J.R. is supported by the Leverhulme Trust. N.L. acknowledges the support of the Swiss National Science Foundation as part of the project ‘Philosophy Beyond Standard Physics’.

References

- [AH17] Felipe A Asenjo and Sergio A Hojman. Do electromagnetic waves always propagate along null geodesics? *Classical and Quantum Gravity*, 34(20):205011, sep 2017.
- [AMCK⁺16] P. Aleahmad, H. Moya Cessa, I. Kaminer, M. Segev, and Christodoulides D. N. Dynamics of accelerating Bessel solutions of Maxwell’s equations. *Journal of the Optical Society of America A*, 33(10):2047, 2016.
- [And15] Sibylle Anderl. Astronomy and astrophysics in the philosophy of science. *arXiv preprint arXiv:1510.03284*, 2015.
- [BB68] J. M. Barnothy and M. F. Barnothy. The relationship between Seyfert galaxies, *n*-type galaxies and quasi-stellar objects. *Astrophysical Letters*, 2:21–26, 1968.
- [BB79] M. V. Berry and N. L. Balazs. Nonspreading wave packets. *American Journal of Physics*, 47(3):264, 1979.
- [BNKS14] Rivka Bekenstein, Jonathan Nemirovsky, Ido Kaminer, and Mordechai Segev. Shape-preserving accelerating electromagnetic wave packets in curved space. *Physical Review X*, 4(1):011038, 2014.
- [Bri60] Léon Brillouin. *Wave Propagation and Group Velocity*. Academic Press, 1960.
- [BSDC08] John Broky, Georgios A Siviloglou, Aristide Dogariu, and Demetrios N Christodoulides. Self-healing properties of optical airy beams. *Optics express*, 16(17):12880–12891, 2008.

¹⁴For a general introduction to notions of reflective equilibrium, see [Dan20]. What we have in mind in this article is akin (we take it) to the iterative revision of the concept of temperature as discussed by Chang [Cha04], but in this case the target phenomena are (of course) electromagnetic signals.

- [But07] Jeremy Butterfield. Reconsidering relativistic causality. *International Studies in the Philosophy of Science*, 21(3):295–328, 2007.
- [Cao01] Tian Yu Cao. Prerequisites for a consistent framework of quantum gravity. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics*, 32(2):181–204, 2001. Spacetime, Fields and Understanding: Perspectives on Quantum Field.
- [CE13] I. D. Chremmos and N. K. Efremidis. Nonparaxial accelerating Bessel-like beams. *Phys. Rev. A*, 88:063816, 2013.
- [Cha04] Hasok Chang. *Inventing Temperature: Measurement and Scientific Progress*. New York, US: OUP Usa, 2004.
- [Cle02] Carol E Cleland. Methodological and epistemic differences between historical science and experimental science. *Philosophy of science*, 69(3):474–496, 2002.
- [CRNW10] A. Chong, W. H. Renninger, Christodoulides D. N., and Wise F. W. Airy–Bessel wave packets as versatile linear light bullets. *Nature Photonics*, 4:103, 2010.
- [Cur18] Adrian Currie. *Rock, Bone, and Ruin: An Optimist’s Guide to the Historical Sciences*. Life and Mind: Philosophical Issues in Biology and Psychology. MIT Press, 2018.
- [Dan20] Norman Daniels. Reflective Equilibrium. In Edward N. Zalta, editor, *The Stanford Encyclopedia of Philosophy*. Metaphysics Research Lab, Stanford University, Summer 2020 edition, 2020.
- [ECSC19] Nikolaos K Efremidis, Zhigang Chen, Mordechai Segev, and Demetrios N Christodoulides. Airy beams and accelerating waves: an overview of recent advances. *Optica*, 6(5):686–701, 2019.
- [Ein05] A. Einstein. Zur Elektrodynamik bewegter Körper. *Annalen der Physik*, 322(10):891–921, January 1905.
- [Ein54] Albert Einstein. *Ideas and Opinions*. Bonanza Books, 1954.
- [GM56] M. Gell-Mann. The interpretation of the new particles as displaced charge multiplets. *Il Nuovo Cimento (1955-1965)*, 4(2):848–866, 1956.
- [GMCB98] J.C. Garrison, M.W. Mitchell, R.Y. Chiao, and E.L. Bolda. Superluminal signals: causal loop paradoxes revisited. *Physics Letters A*, 245(1):19–25, 1998.
- [Hac11] S. Hacyan. Relativistic accelerating electromagnetic waves. *Journal of Optics*, 13:105710, 2011.
- [KBNS12] Ido Kaminer, Rivka Bekenstein, Jonathan Nemirovsky, and Mordechai Segev. Non-diffracting accelerating wave packets of Maxwell’s equations. *Physical Review Letters*, 108:163901, 2012.
- [KH10] Y Kaganovsky and E Heyman. Wave analysis of airy beams. *Optics Express*, 18(8):8440–8452, 2010.
- [Lad01] James Ladyman. *Understanding Philosophy of Science*. Routledge, 2001.
- [Lak74] Imre Lakatos. The role of crucial experiments in science. *Studies in History and Philosophy of Science Part A*, 4(4):309–325, 1974.
- [Lek09] J. Lekner. Airy wavepacket solutions of the schrödinger equation. *Eur. J. Phys.*, 30:L43, 2009.
- [LR21] Niels Linnemann and James Read. Comment on ‘Do electromagnetic waves always propagate along null geodesics?’. *Classical and Quantum Gravity*, 38(23):238001, dec 2021.
- [McD18] Kirk T. McDonald. Slepian’s faster-than-light wave. 2018.

- [NEC⁺18] J. Nichols, Tegan Emerson, L. Cattell, Serim Park, Andrey Kanaev, Frank Bucholtz, Abbie Watnik, T. Doster, and G. Rohde. Transport-based model for turbulence-corrupted imagery. *Applied Optics*, 57:4524, 06 2018.
- [NNB22] J. M. Nichols, D. V. Nickel, and F. Bucholtz. Vector beam bending via a polarization gradient. *Opt. Express*, 30(21):38907–38929, Oct 2022.
- [PBBS18] Anatoly Patsyk, Miguel A Bandres, Rivka Bekenstein, and Mordechai Segev. Observation of accelerating wave packets in curved space. *Physical Review X*, 8(1):011001, 2018.
- [PCP⁺16] P. Peng, B. Chen, X. Peng, M. Zhou, L. Zhang, D. Li, and D. Deng. Self-accelerating airy-ince-gaussian and airy-helical-ince-gaussian light bullets in free space. *Optics Express*, 24(17):18973, 2016.
- [PN98] D. Paganin and K. A. Nugent. Noninterferometric phase imaging with partially coherent light. *Phys. Rev. Lett.*, 80:2586–2589, Mar 1998.
- [PSAT10] D. G. Papazoglou, S. Suntsov, D. Abdollahpour, and S. Tzortzakis. Tunable intense airy beams and tailored femtosecond laser filaments. *Phys. Rev. A*, 81:061807, Jun 2010.
- [SBDC07] G. A. Siviloglou, J. Broky, A. Dogariu, and D. N. Christodoulides. Observation of accelerating airy beams. *Phys. Rev. Lett.*, 99:213901, 2007.
- [SBDC08] G. A. Siviloglou, J. Broky, A. Dogariu, and D. N. Christodoulides. Ballistic dynamics of airy beams. *Opt. Lett.*, 33(3):207–209, Feb 2008.
- [SC07] G. A. Siviloglou and D. N. Christodoulides. Accelerating finite energy airy beams. *Optics Letters*, 32:979, 2007.
- [SGN03] Michael D. Stenner, Daniel J. Gauthier, and Mark A. Neifeld. The speed of information in a ‘fast-light’ optical medium. *Nature*, 425(6959):695–698, 2003.
- [Sho03] G. M. Shore. Causality and superluminal light. In *Time and Matter: An International Colloquium on the Science of Time*, pages 45–66, 2 2003.
- [Sle49] J. Slepian. Waves of all velocities. *Electrical Engineering*, pages 1080–1081, 1949.
- [YSA10] Berna Yalizay, Burak Soylu, and Selcuk Akturk. Optical element for generation of accelerating airy beams. *J. Opt. Soc. Am. A*, 27(10):2344–2346, Oct 2010.
- [ZBH13] W.-P. Zhong, M. Belić, and T. Huang. Three-dimensional finite-energy airy self-accelerating parabolic-cylinder light bullets. *Phys. Rev. A*, 88:033824, 2013.