

# On the geometric trinity of gravity, non-relativistic limits, and Maxwell gravitation

Eleanor March\*, William J. Wolf† & James Read‡

## Abstract

We show that the dynamical common core of the recently-discovered non-relativistic geometric trinity of gravity is Maxwell gravitation. Moreover, we explain why no analogous distinct dynamical common core exists in the case of the better-known relativistic geometric trinity of gravity.

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Geometric trinitities</b>	<b>5</b>
2.1	The relativistic geometric trinity . . . . .	5
2.2	The non-relativistic geometric trinity . . . . .	7
<b>3</b>	<b>Maxwell gravitation and the non-relativistic trinity</b>	<b>9</b>
<b>4</b>	<b>GR as the relativistic common core</b>	<b>15</b>
<b>5</b>	<b>Conclusions</b>	<b>18</b>

## 1 Introduction

*Inter alia*, the following questions surely count as mainstream in contemporary philosophy of spacetime physics:

1. What is the ‘correct’ spacetime setting for Newtonian gravity, especially in light of Newton’s Corollary VI?<sup>1</sup> (On this topic, see e.g. Dewar (2018), Knox (2014), Teh (2018), Wallace (2020), and Weatherall (2016, 2018).)

---

\*Faculty of Philosophy, University of Oxford, UK. [eleanor.march@philosophy.ox.ac.uk](mailto:eleanor.march@philosophy.ox.ac.uk)

†Faculty of Philosophy, University of Oxford, UK. [william.wolf@philosophy.ox.ac.uk](mailto:william.wolf@philosophy.ox.ac.uk)

‡Faculty of Philosophy, University of Oxford, UK. [james.read@philosophy.ox.ac.uk](mailto:james.read@philosophy.ox.ac.uk)

<sup>1</sup>Recall that Newton’s Corollary VI reads as follows: “If bodies are moving in any way whatsoever with respect to one another and are urged by equal accelerative forces along parallel lines, they will all continue to move with respect to one another in the same way as they would if they were not acted on by those forces.” (Newton 2014, p. 99).

2. Are there spacetime theories which are in some sense or other ‘equivalent’ to general relativity, and what would be the philosophical significance of such theories, were they to exist? (On this topic, see e.g. Bain (2006), Duerr and Read (2023), Knox (2011), Rosenstock, Barrett, and Weatherall (2015), Wolf and Read (2023), and Wolf, Sanchioni, and Read (2023).)
3. How is one to take the non-relativistic limit of general relativity, and what is the resulting theory? (On this topic, see e.g. Fletcher (2019) and Malament (1986).)

Until now, discussions of these questions have, broadly speaking, been isolated from one another. Our purpose in this article is to show that these questions (and answers to said questions) are in fact related to one another in intimate and significant ways.

To explain what we mean here, begin with question (1). (For the time being a qualitative account will suffice; the mathematics to substantiate the claims made here will follow later in this article.) Typically, Newtonian gravitation theory (NGT) in its potential-based formulation *chez* Laplace and Poisson is taken to be set in a flat spacetime; gravitational effects in this spacetime are encoded in the gravitation potential which leads to test bodies not traversing geodesics of the flat, compatible connection. This being said, NGT has a hidden symmetry (sometimes referred to as ‘Trautman symmetry’ (Teh 2018)): if one (a) subjects *all* material bodies to an additional constant gravitational field, and (b) changes one’s standard of straightness (i.e., one’s derivative operator) to compensate for this, then in fact no physical change ensues. (This is related to Newton’s Corollary VI, as we will explain below; cf. Read and Teh (2022).) When one moves to a new formalism purged of this additional symmetry,<sup>2</sup> one arrives at the structure of Newton-Cartan theory (NCT): a non-relativistic spacetime theory in which gravitational effects are—just as in the case of general relativity (GR)—manifestations of spacetime curvature.

This much is well-known. But there remains some ambiguity in the literature as to how NGT relates to another spacetime theory known as ‘Maxwell gravitation’ (MG), also developed in light of Newton’s Corollary VI. (On this theory, see Chen (2023), Dewar (2018), March (2023a,b), and Saunders (2013).) Moreover, recently NGT has been shown to in fact admit of an interpretation whereby it is a theory with a *torsionful* geometry (Read and Teh 2018; Schwartz 2023), in the sense that the gravitational potential can be associated with the torsion of the ‘mass gauge field’ which arises when one gauges the Bargmann algebra (for more on the mass gauge field, see Andringa et al. (2011), Read and Teh (2018), Teh (2018), and Wolf, Read, and Teh (2023)). We shall refer to this torsional interpretation of NGT as TENC, or the ‘teleparallel equivalent of Newton-Cartan theory’, for reasons which will become apparent shortly. Even less well-known (indeed, we might say, almost unknown!) is that both TENC and NCT are equivalent to an alternative non-relativistic theory, re-

---

<sup>2</sup>Here, we set aside the differences between what are known as ‘reduction’ and ‘internal sophistication’ about symmetries: see Dewar (2018) and Martens and Read (2021).




Figure 1: Maxwell gravitation as the common core of the non-relativistic geometric trinity of gravity.

cently dubbed the ‘symmetric teleparallel equivalent of Newton-Cartan theory’ (STENC), in which gravitational effects are manifestations neither of curvature (as in NCT) nor of torsion (as in TENC), but of spacetime non-metricity.<sup>3</sup> Here, we demonstrate that these pieces fit together in the following way: NCT, TENC and STENC constitute a ‘non-relativistic geometry trinity’; the structure common to all said theories (the ‘common core’, in the sense of Le Bihan and Read (2018)) *just is* the structure of MG (see Figure 1).<sup>4</sup>

Already, this illuminates quite substantially the connections between these four non-relativistic theories of spacetime and gravity. And yet, that is only the beginning of the story. Taking now together questions (2) and (3) in our above list, it is becoming increasingly well-known to philosophers of physics that there exists a ‘geometric trinity’ of relativistic theories of gravitation, of which GR constitutes but one node (see e.g. Jiménez, Heisenberg, and Koivisto (2019) for a recent review in the physics literature). The other two nodes are ‘teleparallel gravity’ (TEGR), in which gravitational effects are a manifestation of exclusively spacetime torsion, and ‘symmetric teleparallel gravity’ (STEGR), in which gravitational effects are a manifestation of exclusively spacetime non-metricity. Recently, Wolf, Read, and Vigneron (2023) have shown that the above-discussed non-relativistic trinity (*sans* any mention of MG) is indeed the non-relativistic limit (in the sense of a  $1/c$  expansion *à la* Schwartz (2023)) of this relativistic geometry trinity. More specifically, the torsional non-relativistic theory TENC emerges when one takes the non-relativistic limit of the torsional relativistic theory TEGR, while the non-metric non-relativistic theory STENC emerges as the non-relativistic limit of non-metric relativistic theory STEGR. The web of connections is, therefore, as presented in Figure 2 (in that figure, for clarity, we omit MG).

What we add to this discussion in the present paper is an answer to the

<sup>3</sup>To remind the reader: ‘curvature’ quantifies the extent to which parallel transport of a vector along a closed loop doesn’t preserved angles; ‘torsion’ quantifies the extent to which parallel transport in two directions doesn’t commute; ‘non-metricity’ quantifies the extent to which parallel transport of a vector along a closed loop doesn’t preserve the length of that vector. For further background, see e.g. Hehl et al. (1995).

<sup>4</sup>The same notion of a common core is also discussed in e.g. De Haro and Butterfield (2021).




Figure 2: The relativistic geometric trinity and the non-relativistic geometric trinity as its non-relativistic limit.

following question: does there exist a ‘common core’ of the relativistic geometric trinity in the same sense that MG is the common core of the non-relativistic trinity, and if so is it the case that MG is the non-relativistic limit of said relativistic common core? Whilst we address both parts of this question, we should be clear that our answer to the first part is partly in the negative: there is *not* a dynamically distinct common core of the relativistic geometric trinity in the same sense that MG is the dynamically distinct common core of the non-relativistic trinity.

To elaborate: we concur with the recent verdict of Wolf, Sanchioni, and Read (2023), who argue that the common core of all three of GR, TEGR, and STGR simply is just GR. This is because one can always use the metric to build GR’s defining affine structure, as the Levi-Civita connection is the *unique* torsion-free, metric compatible derivative operator; using this derivative operator one can then write down the Einstein equation as usual. In the non-relativistic case there is again a common core—this time, its kinematics include what has been dubbed in the recent philosophical literature a ‘standard of rotation’ (see Weatherall (2018))—; however—and quite differently to the relativistic case!—that common core leads to dynamics which are *distinct* from those of the the geometric trinity from which they arise. Later in this article, we will explain exactly how this asymmetry between the relativistic and non-relativistic cases comes about.

To summarise, then, in this article we (a) identify MG as the dynamical common core of the recently-discovered non-relativistic geometric trinity of gravity, and (b) explain how it can be that no analogous common core exists in the case of the relativistic geometric trinity of gravity. In so doing, we (i) clarify questions in (1) regarding the ‘correct’ spacetime setting for Newtonian gravity, (ii) connect that entire literature up to the geometric trinity of gravity and its Newtonian limit, which has also aroused recent philosophical interest.

More specifically, the structure of the article is this. In §2, we remind the reader of the mathematical details of both the relativistic geometric trinity and the non-relativistic geometric trinity. In §3, we present MG as the common core

of the non-relativistic trinity, and connect our discussion to that of the ‘correct’ spacetime setting for Newtonian gravity. In §4, we address the matter of the existence (or otherwise) of a relativistic common core. We close in §5.

## 2 Geometric trinities

In this section, we recall the mathematical details underlying the existence of the relativistic geometric trinity of gravitation theories (§2.1) and the non-relativistic geometric trinity of gravitation theories (§2.2).

### 2.1 The relativistic geometric trinity

The ‘geometric trinity’ of gravity refers to a family of three relativistic theories of gravitation: general relativity (GR), the ‘teleparallel equivalent to general relativity’ (TEGR), and the ‘symmetric teleparallel equivalent to general relativity’ (STTEGR). These theories are all ‘equivalent’ to each other in the sense that they share equivalent dynamical equations of motion, but distinct in the sense that these shared dynamics result from entirely different geometric degrees of freedom that manifest in each respective theory (see e.g. Capozziello, De Falco, and Ferrara (2022), Heisenberg (2019), Jiménez, Heisenberg, and Koivisto (2019), and Wolf, Read, and Vigneron (2023)).

Kinematical possibilities of general relativity are typically presented as tuples of the form  $\langle M, g_{ab}, \Phi \rangle$ , where  $M$  is a four-dimensional differentiable manifold,  $g_{ab}$  is a Lorentzian metric field on  $M$ , and  $\Phi$  represents material fields. The dynamical possibilities of the theory are encoded by the Einstein equation, which governs the behavior of these spacetime and material fields. However, the geometric degrees of freedom responsible for sourcing the dynamics of the respective theories in the geometric trinity are properties of the affine connection. We will thus take GR to be a theory given by models of the form  $\langle M, g_{ab}, \overset{c}{\nabla}, \Phi \rangle$ , where  $\overset{c}{\nabla}$  refers to the familiar Levi-Civita derivative operator with non-vanishing *curvature*. Typically,  $\overset{c}{\nabla}$  is not included explicitly in the models of GR, for it is fixed uniquely by  $g_{ab}$ .

Spacetime curvature is defined in the following way:

$$R^a{}_{bcd}\xi^b := -2\nabla_{[c}\nabla_{d]}\xi^a, \quad (1)$$

where  $\xi^a$  is a smooth vector field. However, curvature is not the only geometric property that a connection can manifest. An affine connection can also possess *torsion* or *non-metricity*. The torsion tensor is given by

$$T^c{}_{ab}\nabla_c\alpha := 2\nabla_{[a}\nabla_{b]}\alpha, \quad (2)$$

where  $\alpha$  is a smooth scalar field; torsion thereby encodes the antisymmetry of a connection. Non-metricity is given by the non-vanishing of the covariant derivative of the metric tensor

$$Q_{abc} := \nabla_a g_{bc}. \quad (3)$$

Heuristically, curvature measures the rotation of a vector when it is parallel transported along a closed curve, torsion measures of the non-closure of the parallelogram formed by two vectors being parallel transported along each other, and non-metricity measures how the length of a vector changes when parallel transported (see e.g. Figure 1 in Jiménez, Heisenberg, and Koivisto (2019) or Hehl et al. 1995).

The Levi-Civita connection of GR is unique in the sense that it is the unique derivative operator which is both torsion-free (i.e.  $T^a_{bc} = 0$ ) and metric-compatible (i.e.  $Q_{abc} = 0$ ), but with generically non-vanishing curvature (i.e.  $R^a_{bcd} \neq 0$ ). However, in order to build a viable relativistic spacetime theory, it is not necessary to use  $\overset{c}{\nabla}$ . Indeed, one can decompose a general affine connection as

$$\nabla = (\overset{c}{\nabla}, K^a_{bc} + L^a_{bc}), \quad (4)$$

where  $K^a_{bc}$  is known as the ‘contorsion tensor’ and  $L^a_{bc}$  is known as the ‘distorsion tensor’ (here, we use the notation of Malament (2012, p. 53)). The contorsion tensor can be understood as the difference tensor between the Levi-Civita connection and the torsionful (but flat and metric-compatible) connection of TEGR. The distorsion tensor can be understood as the difference tensor between the Levi-Civita connection and non-metric (but flat and torsionless) connection of STEGR.

If—as above—we take GR to be a theory with kinematical possibilities of the form  $\langle M, g, \overset{c}{\nabla}, \Phi \rangle$ , then TEGR can be taken to be a theory with kinematical possibilities given by  $\langle M, g, \overset{t}{\nabla}, \Phi \rangle$ , where  $\overset{t}{\nabla} = (\overset{c}{\nabla}, K^a_{bc})$  refers to the TEGR connection with vanishing curvature and non-metricity, but non-vanishing torsion. Likewise, STEGR is a theory with kinematical possibilities given by  $\langle M, g, \overset{n}{\nabla}, \Phi \rangle$ , where  $\overset{n}{\nabla} = (\overset{c}{\nabla}, L^a_{bc})$  refers to the STEGR connection with vanishing curvature and torsion, but non-vanishing non-metricity.

One can use (4) as a dictionary by which to translate between these theories. That is, one can rewrite the geometric objects of interest in one theory in terms of the geometric objects of one of the other trinity theories, and thereby witness their equivalence. For example, one can take the curvature tensor for a generic affine connection and use (4) to relate the curvature of the Levi-Civita connection to that of the TEGR connection (and the associated contorsion tensor), or to that of the STEGR connection (and the associated distorsion tensor). One then finds that the geometric scalar quantities have the following relationship:

$$-R = T + B_T = Q + B_Q, \quad (5)$$

where  $R$  is the curvature scalar,  $T$  is the torsion scalar,  $Q$  is the non-metricity scalar, and  $B_{T/Q}$  refers to boundary terms of the respective theories (Jiménez, Heisenberg, and Koivisto 2019). This also illustrates that these theories are dynamically equivalent, as the Lagrangian expressions for all of these theories can be written using the geometric scalars (in the case of GR, recall the Einstein-Hilbert action). Upon utilizing standard variational procedures, the boundary

terms that arise in (5) vanish, resulting in the Einstein equation for all theories (but of course expressed in their particular geometric languages).<sup>5</sup>

## 2.2 The non-relativistic geometric trinity

It was shown recently by Wolf, Read, and Vigneron (2023) that there is a non-relativistic analogue of the geometric trinity, whereby standard Newtonian gravity can likewise be reconceptualised and/or reformulated as a theory of curvature, torsion, or non-metricity. The three nodes of the non-relativistic geometric trinity of gravity are ‘Newton-Cartan theory’ (NCT), the ‘teleparallel equivalent of Newton-Cartan theory’ (TENC), and the ‘symmetric teleparallel equivalent of Newton-Cartan theory’ (STENC).

Following the presentation of the relativistic theories above, we take NCT to be a theory with kinematical possibilities of the form  $\langle M, t_a, h^{ab}, \overset{c}{\nabla}, \Phi \rangle$ . As before,  $M$  is a four-dimensional differentiable manifold,  $\Phi$  represents material fields, and  $\overset{c}{\nabla}$  is a torsion-free and compatible (now in the sense that  $\nabla_a t_b = \nabla_a h^{bc} = 0$ ) derivative operator with non-vanishing curvature. However, there are some important differences:

1. The metrical structure of non-relativistic theories is notably different from that of relativistic theories, because  $t_a$  and  $h^{ab}$  refer to degenerate temporal and spatial metrics on  $M$ : see Malament (2012, Ch. 4).<sup>6</sup> Metric compatibility applies separately to both metrics; in addition,  $t_a$  and  $h^{ab}$  are orthogonal to each other, so that  $t_a h^{ab} = 0$ . Loosely speaking,  $t_a$  is supposed to represent Newtonian absolute time, and  $h^{bc}$  is supposed to represent Newtonian absolute space.
2. The dynamical possibilities for NCT are encoded in the ‘geometrised Poisson equation’:

$$R_{ab} = 4\pi\rho t_a t_b, \tag{6}$$

where  $R_{ab}$  is the Ricci curvature of the NCT connection  $\overset{c}{\nabla}$ ; moreover, one typically includes the following curvature conditions in one’s presentation of NCT (we will discuss these curvature conditions more later in the article<sup>7</sup>):

$$R^a{}_b{}^c{}_d = R^c{}_d{}^a{}_b, \tag{7}$$

$$R^{ab}{}_{cd} = 0. \tag{8}$$

---

<sup>5</sup>For more on the significance of these boundary terms, see Wolf and Read (2023) for philosophical discussion on concerning their implications for theory equivalence and theory structure and see Oshita and Wu (2017) for further physics discussion.

<sup>6</sup>Through this article, we assume temporal orientability, in the sense of Malament (2012, ch. 4).

<sup>7</sup>See also Malament (2012) and Teh (2018) for further discussion of the physical significance and meaning of these conditions.

TENC is a theory given by models of the form  $\langle M, t_a, h^{ab}, \overset{t}{\nabla}, \Phi \rangle$ , where  $\overset{t}{\nabla}$  again is a flat, metric-compatible derivative operator with (in a certain quite non-obvious sense which we will explain!) non-vanishing torsion. One can translate between NCT and TENC by introducing the difference tensor  $\overset{t}{\nabla} = (\overset{c}{\nabla}, t_b t_c \overset{t}{\nabla}^a \phi)$ , where  $\phi$  is a scalar field representing the gravitational potential. This allows one to cast the dynamics of this theory in its familiar form given by the standard Poisson equation of NGT:

$$\overset{t}{\nabla}_a \overset{t}{\nabla}^a \phi = 4\pi\rho. \quad (9)$$

Although the translation between NCT and NGT (in the form of the the famous ‘geometrisation’ and ‘recovery’ theorems—(see also Malament (2012, ch. 4)) has been known since the work of Trautman (1965), it was much more recently shown by Read and Teh (2018) that standard Newtonian gravity can be understood as the ‘teleparallel equivalent’ of NCT in much the same way that TEGR is the teleparallel equivalent of GR. Here, the gravitational field follows from the torsion of the ‘mass gauge field’  $m_a$ , which is obtained by gauging the Bargmann algebra: one has  $T(M) := dm_a$ ; gauge fixing  $m_a = \phi t_a$  yields (9). In other words, TENC, despite its formal equivalence to NGT, interprets gravity as a result of a torsional connection where this torsion is sourced by the mass gauge field.

Even more recently, STENC has been constructed by Wolf, Read, and Vigneron (2023). This theory is given by models of the form  $\langle M, t_a, h^{ab}, \overset{n}{\nabla}, \Phi \rangle$ , where  $\overset{n}{\nabla}$  is flat and torsion-free but possesses non-vanishing non-metricity. Similarly, one can translate between NCT and STENC by introducing the difference tensor  $\overset{n}{\nabla} = (\overset{c}{\nabla}, v^\alpha Q_{(bc)} - t_b t_c \overset{n}{\nabla}^a \phi)$  and obtain an equivalent equation to (6) and (9) cast now in terms of non-metricity degrees of freedom:

$$\overset{n}{\nabla}_a \overset{n}{\nabla}^a \phi = 4\pi\rho. \quad (10)$$

As with TENC above, one obtains an equation formally equivalent to NGT, but now the gravitational field is interpreted as resulting from a connection that is flat and torsionless, but that encodes non-metricity of the form:

$$\begin{aligned} \overset{n}{\nabla}_a t_b &= Q_{ab}, \\ \overset{n}{\nabla}_a h^{bc} &= 2v^{(b} Q_{ad} h^{c)d}, \end{aligned} \quad (11)$$

where  $v^a$  is a timelike vector field such that  $v^a t_a = 1$ . In order to obtain this formal equivalence with NGT, one must also (as before with TENC and gauge fixing the form of the torsion), gauge fix the form of the STENC non-metricity (see Wolf, Read, and Vigneron (2023, Sect. 4.3) for details).

This ‘non-relativistic geometric trinity’ mirrors the more familiar relativistic geometric trinity in that one can present three gravitational theories that are



dynamically equivalent to familiar Newtonian gravitational theory, formulated in the geometric languages of curvature, torsion, and non-metricity. While in the relativistic case this is apparent at the level of the action, NCT, TENC, and STENC by contrast cannot be formulated using an action principle (for the reasons underlying this, see Hansen, Hartong, and Obers (2019)), so we can only demonstrate their equivalence via Trautman-style geometrisation and recovery theorems.<sup>8</sup> However, the non-relativistic trinity bears another important relationship to the relativistic trinity. All of the theories in the non-relativistic trinity can be obtained by taking an appropriate non-relativistic limit (typically in terms of a  $1/c$  expansion in the style of Schwartz (2023)). That is, NCT is the non-relativistic limit of GR (Künzle 1976), TENC is the non-relativistic limit of TEGR (Schwartz 2023), and STENC is the non-relativistic limit of STEGR (Wolf, Read, and Vigneron (2023)). This completes all the legs of Figure 2.

### 3 Maxwell gravitation and the non-relativistic trinity

As emphasised in the previous section, the three nodes of each of the relativistic and non-relativistic geometric trinities are all empirically equivalent theories, which nevertheless appear to disagree fundamentally on the geometrical structure which they attribute to the world. For instance, according to GR the gravitational behaviour of matter is to be understood as a manifestation of spacetime curvature, whereas according to TEGR and STEGR spacetime is necessarily flat. This means that the relativistic and non-relativistic theories present a case of strong underdetermination—distinct theories between which no possible evidence could be expected to decide.<sup>9</sup>

Faced with such cases of strong underdetermination, philosophers have suggested several approaches to dealing with the problem (on this see e.g. Le Bihan and Read (2018)). Famously, one of these is the common core approach. The common core approach advocates identifying the invariant kinematical structure of the theories, and then showing that this structure is sufficient to formulate a distinct, ontologically viable theory in its own right; one which, moreover, retains the empirical content of the original theories. Moving to this new interpretative framework alongside a judicious invocation of Occamist norms (on which see Dasgupta (2016)) then allows one to ‘break’ the underdetermination by interpreting the theories in such a way that they completely agree on the structure they attribute to the world. The aim of this section is to show that in the case of the non-relativistic geometric trinity, such a common core theory exists, and it is a theory known as ‘Maxwell gravitation’: a theory which has quite independently attracted philosophical interest (for reasons to do with (1)

<sup>8</sup>Off-shell non-relativistic equivalence would require recourse to the ‘Type II’ versions of these theories, which we discuss in §5.

<sup>9</sup>Knox (2011) argues on the basis of her ‘spacetime functionalism’ that there is in fact no strong underdetermination here; see (Mulder and Read 2023; Wolf, Sanchioni, and Read 2023) for responses, and §4 for further discussion in this article.

as presented in the introduction).

To do so, we begin by recalling some facts about Maxwellian spacetime. This is a structure  $\langle M, t_a, h^{ab}, \circ \rangle$ , where  $t_a, h^{ab}$  are orthogonal temporal and spatial metrics as introduced in the previous section, and  $\circ$  is a standard of rotation compatible with  $t_a$  and  $h^{ab}$ . This was introduced originally by Weatherall (2018): if  $t_a, h^{ab}$  are compatible temporal and spatial metrics on  $M$ , then a standard of rotation  $\circ$  compatible with  $t_a$  and  $h^{ab}$  is a map from smooth vector fields  $\xi^a$  on  $M$  to smooth, antisymmetric rank-(2,0) tensor fields  $\circ^b \xi^a$  on  $M$ , such that

1.  $\circ$  commutes with addition of smooth vector fields;
2. Given any smooth vector field  $\xi^a$  and smooth scalar field  $\alpha$ ,  $\circ^a (\alpha \xi^b) = \alpha \circ^a \xi^b + \xi^{[b} d^a] \alpha$ ;
3.  $\circ$  commutes with index substitution;
4. Given any smooth vector field  $\xi^a$ , if  $d_a(\xi^n t_n) = 0$  then  $\circ^a \xi^b$  is spacelike in both indices; and
5. Given any smooth spacelike vector field  $\sigma^a$ ,  $\circ^a \sigma^b = D^{[a} \sigma^{b]}$ ,

where  $D$  is the unique Levi-Civita connection induced by  $h^{ab}$  on each spacelike hypersurface. We will say that a connection and a standard of rotation are *compatible* iff  $\circ^a \eta^b = \nabla^{[a} \eta^{b]}$  for all vector fields  $\eta^a$  on  $M$ . It follows that any compatible, torsion-free connection  $\nabla$  on  $M$  determines a unique compatible standard of rotation—namely, the map  $\circ: \eta^a \rightarrow \nabla^{[a} \eta^{b]}$  (Weatherall 2018).

However, in light of the discussion of the previous section, this invites a natural further question: are there non-metric or torsionful connections which are *also* associated with metric compatible standards of rotation in the above sense? It turns out that the answer to this question is ‘yes’, and a characterisation of such connections is given by the following two propositions:

**Proposition 1.** *Let  $\langle M, t_a, h^{ab}, \nabla \rangle$  be a non-relativistic spacetime, where  $t_a$  and  $h^{ab}$  are compatible, and where  $\nabla_a t_b = Q_{ab}$  and  $\nabla_a h^{bc} = Q_a{}^{bc}$ . Then the map  $\circ: \eta^a \rightarrow \nabla^{[a} \eta^{b]}$  is a standard of rotation compatible with  $t_a$  and  $h^{ab}$  iff  $h^{an} Q_{nb} = h^{n[a} Q_n{}^{b]c} = 0$ .*

*Proof.* First, suppose that  $h^{an} Q_{nb} = h^{n[a} Q_n{}^{b]c} = 0$ . That  $\circ$  satisfies conditions (1)-(3) is immediate from properties of derivative operators. For condition (4), note that if  $d_a(\eta^n t_n) = 0$  we have

$$0 = \nabla_a(\eta^n t_n) = t_n \nabla_a \eta^n + \eta^n Q_{na}, \quad (12)$$

so that

$$\begin{aligned}
t_n(h^{m[n}\nabla_m\eta^{a]}) &= -\frac{1}{2}h^{ma}t_n\nabla_m\eta^n \\
&= \frac{1}{2}h^{ma}\eta^n Q_{mn} \\
&= 0 \\
&= t_n(h^{m[a}\nabla_m\eta^{n]})
\end{aligned}$$

and  $\circlearrowleft^a\eta^b$  is spacelike in both indices. Finally, consider condition (5). We know that  $\nabla^{[a}h^{b]c} = h^{n[a}Q_n{}^{b]c} = 0$ . So let  $\xi^a$  be a unit timelike vector field on  $M$ ,  $\hat{h}_{ab}$  the spatial metric relative to  $\xi^a$ ,<sup>10</sup> and  $D$  the unique spatial derivative operator such that  $D_a h^{bc} = 0$ . Then for any spacelike vector field  $\sigma^a$  on  $M$ ,

$$h^{n[a}D_n\sigma^{b]} = h^{n[a}\hat{h}_{nm}\hat{h}^b]_r\nabla^m\sigma^r = h^{n[a}\nabla_n\sigma^{b]}, \quad (13)$$

where the first equality follows from the proof of (Weatherall 2018, Proposition 2), and we have used the fact that since  $h^{an}Q_{nb} = 0$ ,  $\nabla^a(t_n\eta^n) = t_n\nabla^a\eta^n$  for any smooth vector field  $\eta^a$  on  $M$ .

Conversely, suppose that the map  $\circlearrowleft: \eta^a \rightarrow \nabla^{[a}\eta^{b]}$  is compatible with  $t_a$  and  $h^{ab}$ . Then we must have for all smooth vector fields  $\eta^a$  on  $M$  that if  $d_a(\eta^n t_n) = 0$ ,  $\circlearrowleft^a\eta^b$  is spacelike in both indices. If  $d_a(\eta^n t_n) = 0$  then (12) holds with respect to  $\eta^a$  so that  $t_n(h^{m[n}\nabla_m\eta^{a]}) = 1/2h^{ma}Q_{mn}\eta^n = 0$ . But this can only be the case for arbitrary  $\eta^a$  if  $h^{an}Q_{nm} = 0$ . Moreover, given condition (5) we must then also have that the equality (13) holds with respect to  $\nabla$ . Now, we know that  $h^{n[a}D_n\sigma^{b]} = h^{n[a}\hat{h}_{nm}\hat{h}^b]_r\nabla'^m\sigma^r$  for any  $\nabla'$  such that  $\nabla'^a h^{bc} = 0$ ; in particular, this is the case for  $\nabla' = (\nabla, -1/2\hat{h}_{bn}\hat{h}_{cm}(h^{ar}Q_r{}^{nm} - h^{nr}Q_r{}^{am} - h^{mr}Q_r{}^{na}))$ . Thus

$$\begin{aligned}
0 &= h^{n[a}\hat{h}_{nm}\hat{h}^b]_r h^{ms}\hat{h}_{st}\hat{h}_{vu}(h^{rw}Q_w{}^{tu} - h^{tw}Q_w{}^{ru} - h^{uw}Q_w{}^{tr})\sigma^v \\
&= \hat{h}^{[a}{}_{t}\hat{h}^b]_r\hat{h}_{vu}(h^{rw}Q_w{}^{tu} - h^{tw}Q_w{}^{ru} - h^{uw}Q_w{}^{tr})\sigma^v \\
&= 2\delta^{[a}{}_{t}\delta^{b]}_r\hat{h}_{vu}h^{w[r}Q_w{}^{t]u}\sigma^v \\
&= h^{w[b}Q_w{}^{a]u}\sigma_u
\end{aligned}$$

for any covector  $\sigma_a$ , where we have used that the metrics are orthogonal (so that  $t_n Q^{ab}$  vanishes if  $Q^a{}_b$  does) and that  $Q_a{}^{bc}$  is symmetric in the upper two indices. But this can only be the case for arbitrary  $\sigma_a$  if  $h^{n[a}Q_n{}^{b]c} = 0$ .  $\square$

**Proposition 2.** *Let  $\langle M, t_a, h^{ab}, \nabla \rangle$  be a non-relativistic spacetime, where  $t_a$  and  $h^{ab}$  are compatible, and  $\nabla$  is compatible with the metrics but possibly torsionful. Then the map  $\circlearrowleft: \eta^a \rightarrow \nabla^{[a}\eta^{b]}$  is a standard of rotation compatible with  $t_a$  and  $h^{ab}$  iff  $T^{abc} = 0$ .*

<sup>10</sup>That is, the unique symmetric tensor field on  $M$  such that  $\hat{h}_{an}\xi^n = 0$  and  $h^{an}\hat{h}_{nb} = \delta^a{}_b - t_b\xi^a$ .

*Proof.* First, note that since  $\nabla$  is compatible with  $t_a$  and  $t_a$  is closed,  $t_n T^n{}_{ab} = 0$ . That the map  $\circlearrowleft: \eta^a \rightarrow \nabla^{[a}\eta^{b]}$  satisfies conditions (1)-(3) is again immediate from properties of derivative operators. (4) follows from the fact that  $\nabla$  is compatible with  $t_a$ , using that  $d_a\alpha = \nabla_a\alpha$  for any 0-form field  $\alpha$ . Finally, consider (5). Let  $\xi^a$  be a unit timelike vector field on  $M$ , and  $\hat{h}_{ab}$  the spatial metric relative to  $\xi^a$ . We know that the action of  $D$  on spacelike vector fields is defined as follows:  $D_a\sigma^b = \hat{h}_{an}\hat{h}^b{}_m\nabla'^n\sigma^m$ , where  $\nabla'$  is an arbitrary torsion-free derivative operator such that  $\nabla'^a\hat{h}^{bc} = 0$ . Moreover, we know that  $\nabla = (\nabla', K^a{}_{bc})$  for some such  $\nabla'$ , where  $K^a{}_{bc} = 1/2T^a{}_{bc} + T_{(b}{}^a{}_{c)} + h^{an}t_{(b}f_{c)n}$  is the Newton-Cartan contorsion (Schwartz 2023). Hence

$$\begin{aligned} D^{[a}\sigma^{b]} &= \hat{h}^{[a}{}_n\hat{h}^{b]}{}_m\nabla'^n\sigma^m \\ &= \hat{h}^{[a}{}_n\hat{h}^{b]}{}_m(\nabla^n\sigma^m - h^{nr}K^m{}_{rs}\sigma^s) \\ &= \nabla^{[a}\sigma^{b]} - K^{[ab]}{}_n\sigma^n \\ &= \nabla^{[a}\sigma^{b]} - \frac{1}{2}T^{nab}\sigma_n \end{aligned}$$

for some covector  $\sigma_n$ , where we have used the fact that  $\sigma^a$  is spacelike and  $t_n T^n{}_{ab} = 0$ . Thus if  $T^{abc} = 0$ , then (5) is satisfied. Conversely, if the map  $\circlearrowleft: \eta^a \rightarrow \nabla^{[a}\eta^{b]}$  satisfies (5), it follows that  $T^{abc} = 0$ .  $\square$

Propositions 1 and 2 are tantalising, because they show that non-relativistic affine connections which exhibit either torsion or non-metricity may—under certain conditions—be associated with a compatible standard of rotation, just as with curvature based connections. This raises the prospect that Maxwellian spacetime might be the invariant kinematic structure of the non-relativistic geometric trinity. We isolate the sense in which this is so in the following proposition:

**Proposition 3.** *Consider the triple  $\langle M, t_a, h^{ab} \rangle$  with  $t_a$  and  $h^{ab}$  defined as above. Consider three connections: a curvature based connection  $\overset{c}{\nabla}$ , a non-metricity based connection  $\overset{n}{\nabla}$  with  $\overset{n}{\nabla}_a t_b = Q_{ab}$  and  $\overset{n}{\nabla}_a h^{bc} = Q_a{}^{bc}$ , and a torsion based connection  $\overset{t}{\nabla}$ . Let  $\overset{c}{\nabla} = (\overset{n}{\nabla}, L^a{}_{bc}) = (\overset{t}{\nabla}, K^a{}_{bc})$ , where  $L^a{}_{bc} = 1/2\hat{h}_{bn}\hat{h}_{cm}(h^{ar}Q_r{}^{nm} - h^{nr}Q_r{}^{am} - h^{mr}Q_r{}^{na}) + h^{an}t_{(b}f_{c)n}$  and  $K^a{}_{bc}$  is as in Proposition 2. Then  $\overset{c}{\circlearrowleft}, \overset{n}{\circlearrowleft}, \overset{t}{\circlearrowleft}$  are standards of rotation compatible with the metrics and  $\overset{n}{\circlearrowleft} = \overset{c}{\circlearrowleft} = \overset{t}{\circlearrowleft}$  iff  $h^{an}Q_{nb} = h^{n[a}Q_n{}^{b]c} = 0$ ,  $T^{abc} = 0$ , and  $f_{ab} = t_{[a}\phi_{b]}$  for some  $\phi_a$ .*

*Proof.* That  $\overset{c}{\circlearrowleft}$  is a standard of rotation compatible with the metrics follows immediately from Proposition 1 of Weatherall (2018). The remainder of the proof just involves some calculation. Let  $\eta^a$  be an arbitrary vector field on  $M$ . We know (Proposition 1) that  $\overset{n}{\circlearrowleft}$  is compatible with the metrics iff  $h^{an}Q_{nb} = h^{n[a}Q_n{}^{b]c} = 0$ . So for the first half of the equality, we need to show that if

$h^{an}Q_{nb} = h^{n[a}Q_n{}^{b]c} = 0$  then  $\overset{c}{\nabla} = \overset{n}{\nabla}$  iff  $f_{ab} = t_{[a}\phi_b]$  for some  $\phi_a$ . We have

$$\begin{aligned}\overset{c}{\nabla}^{[a}\eta^{b]} - \overset{n}{\nabla}^{[a}\eta^{b]} &= -L^{[ab]}{}_n \eta^n \\ &= \frac{1}{2}(\hat{h}_{mn}h^{r[a}Q_r{}^{b]m} - t_n f^{[ab]})\eta^n \\ &= -\frac{1}{2}t_n f^{ab})\eta^n \\ &= 0,\end{aligned}$$

where we have used that  $f_{ab}$  is antisymmetric. Since  $t_a$  is nowhere vanishing and  $\eta^a$  is arbitrary, this last term will vanish just in case  $f^{ab} = 0$ . But this is equivalent to the requirement that  $f_{ab} = t_{[a}\phi_b]$  for some covector  $\phi_a$ .

For the second part of the equality, let  $\xi^a$  be a unit timelike vector field, and  $\hat{h}_{ab}$  the spatial metric relative to  $\xi^a$ . Given Proposition 2, we need to show that if  $T^{abc} = 0$ , then  $\overset{c}{\nabla} = \overset{t}{\nabla}$  iff  $f_{ab} = t_{[a}\phi_b]$  for some  $\phi_a$ . Analogously to the first part of the proof

$$\begin{aligned}\overset{c}{\nabla}^{[a}\eta^{b]} - \overset{t}{\nabla}^{[a}\eta^{b]} &= -K^{[ab]}{}_n \eta^n \\ &= -\frac{1}{2}(T_n{}^{ab} + t_n f^{[ab]})\eta^n \\ &= -\frac{1}{2}(\hat{h}_{nm}T^{mab} + t_n f^{ab})\eta^n \\ &= -\frac{1}{2}t_n f^{ab}\eta^n,\end{aligned}$$

which completes the proof by the same reasoning as above.  $\square$

The significance of Proposition 3 stems from the fact that in both STENC and TENC, the homogeneous Trautman conditions (7) and (8) ensure that we are always free to ‘gauge fix’ the non-metricity and torsion so that  $h^{an}Q_{nb} = h^{n[a}Q_n{}^{b]c} = 0$ ,  $T^{abc} = 0$ , and  $f_{ab} = t_{[a}\phi_b]$ . The result is that each model of NCT, and its associated models of STENC and TENC, all pick out a unique Maxwellian spacetime.

Not only this, but the condition (8) guarantees in addition that this Maxwellian spacetime is unique, up to isomorphism, across the entire space of models.<sup>11</sup> So Proposition 3 in fact gives us the stronger result: NCT, TENC, and STENC can all be understood as theories built on the very same Maxwellian spacetime background.

This agreement is striking. It is also to be expected. Stepping back for a minute, all three nodes of the non-relativistic geometric trinity are supposed to be empirically equivalent to standard potential-based Newtonian gravity

<sup>11</sup>This follows since (8) holds with respect to  $\overset{c}{\nabla}$  iff  $\overset{c}{\nabla}$  is rotationally equivalent to some flat compatible torsion-free connection (Malament 2012, proposition 4.2.4; Weatherall 2018, proposition 1), and flat compatible torsion-free connections are unique up to isomorphism.

(i.e. NGT *sans* a torsionful interpretation). *That* theory is one in which spacetime is flat, metric, and (spatiotemporally) torsion-free. Moreover, the dynamics of Newtonian gravitation pick out a privileged class of frames which are non-rotating with respect to that connection.<sup>12</sup> Any theory which is empirically equivalent to Newtonian gravitation must be able to distinguish this same class of frames, which requires agreement on the Maxwellian spacetime structure.

We can see this in more detail by considering the conditions which ensure agreement of the rotation standard. As remarked before, these amount to the homogeneous Trautman conditions. It is well known that only one of these conditions (7) emerges in the non-relativistic limit of GR. The second Trautman condition (8) must be put in by hand as an additional assumption once the non-relativistic limit has been taken.<sup>13</sup> In light of this, a well-known result by Künzle and Ehlers proceeds to generalise the Trautman recovery theorem by dropping the condition (8) (see Malament (2012, Ch. 4) for discussion). The resulting theory, however, is *not* empirically equivalent to standard potential-based Newtonian gravity, but leads to ineliminable coriolis force terms in the recovered models.

To summarise the results of this section so far, we’ve isolated a Maxwellian spacetime structure as the common core of the non-relativistic geometric trinity. We’ve also seen that the constraints which guarantee agreement on the Maxwellian spacetime structure are precisely the constraints which are standardly imposed after taking the non-relativistic limit to ensure empirical equivalence with the potential-based formulation of Newtonian gravity. The next point to note is that this structure is sufficient to formulate the dynamics of Newtonian gravity. This was first done in Dewar (2018), and recently given an ‘intrinsic’ formulation by Chen (2023) and March (2023b); the resulting theory—‘Maxwell gravitation’—has models of the form  $\langle M, t_a, h^{ab}, \circlearrowleft, \Phi \rangle$ . Together with propositions 1, 2, and 3, this substantiates our earlier claim that MG constitutes the dynamical common core of the non-relativistic geometric trinity. It also paves the way for an interpretation of Newtonian gravity on which the structure it attributes to the world is strictly *less* than that of a connection.

This takes us to the connection with Corollary VI and the Trautman symmetry, to which we alluded in Section 1. As articulated by Jacobs (2023), the ‘dynamic shift’ symmetry of potential-based Newtonian gravity *à la* Corollary VI and the Trautman symmetry in which the connection and gravitational potential are altered simultaneously can be understood as being two sides of the same coin: both consequences of the invariance of the Newtonian dynamics under the Maxwell group. But Maxwell transformations produce a linear, time-dependent acceleration of the matter content of the original solution. *Prima facie*, one might think that purging the theory of this symmetry would involve excising the structure needed to make sense of such linear accelerations—to wit,

<sup>12</sup>Here we set aside substantialist-relationalist questions of whether this is a correct prediction of Newtonian theory, on which see e.g. Barbour (2010) and Pooley (2013).

<sup>13</sup>Although one can recover this condition by restricting attention to spacetimes which are, in a certain weak sense, asymptotically flat when taking the non-relativistic limit—on which, see e.g. Malament (2012, Ch. 4.5).

a connection—leaving only the standard of rotation.

In that sense, that MG should be the dynamical common core of the non-relativistic geometric trinity was already suggested by the dynamical symmetries of Newtonian gravity. On the one hand, the irrotational degrees of freedom of the connection were already known to be superfluous to the internal dynamics of the matter distribution. On the other hand, agreement on the rotation standard is necessary for empirical equivalence to standard Newtonian gravity.

What, then, to make of the fact that one can also use Trautman symmetry to motivate the move to NCT? One way to think about this is that while we are always free to *define* a connection from the standard of rotation and matter distribution by coupling the degrees of freedom of the connection to the matter distribution, there is necessarily a certain amount of slack in how this connection is constructed. This is because the projective degrees of freedom of an affine connection far outstrip the degrees of freedom of the matter distribution. Taking up this slack in different places allows us to express Newtonian gravity as a theory of curvature, or torsion, or non-metricity—in some cases, we can even specify the connection uniquely! That we can specify the curvature-based connection uniquely under certain weak conditions on the mass density field is what ensures that  $\overset{c}{\nabla}$ , as well as the rotation standard, is also an invariant of Trautman gauge symmetry. But the fact remains *viz-à-viz* Corollary VI that the full structure of an affine connection is not needed to support the dynamics, and so, if one introduces such a connection, one has to reckon with the fact that—again, necessarily—there are multiple distinct ways of doing so.

## 4 GR as the relativistic common core

Recall that the projective structure of a spacetime theory identifies a certain subset of worldlines the (unparameterised) geodesics; the conformal structure of a given relativistic spacetime theory specifies a lightcone at every spacetime point (see e.g. Matveev and Scholz (2020)). Famously—a result going back to Weyl (1921)—a Lorentzian spacetime is fixed by its associated projective and conformal structure: see Malament (2012, ch. 2); the corresponding existence result was proved by Ehlers, Pirani, and Schild (2012), and is discussed further by Linnemann and Read (2021). In Wolf, Sanchioni, and Read (2023), it was identified that one can move between nodes of the relativistic geometric trinity by modifying projective structure while leaving conformal structure unchanged (for all three theories leave lightcone structure unmodified); in a similar manner (one ultimately irrelevant to our purposes here, but perhaps nevertheless worth pointing out) one can move to non-relativistic theories by ‘widening the lightcone’, thereby changing conformal structure (this constitutes a geometrical way of thinking about taking the non-relativistic limit: see Malament (1986)).

The three nodes of the relativistic geometric trinity then all agree on conformal structure. But the reason for this is that they all agree on metric structure (see §2.1). This means that the invariant kinematical structure of the relativistic geometric trinity is  $\langle M, g_{ab}, \phi \rangle$ . Is this structure sufficient to formulate the

dynamics of the relativistic trinity—i.e. a dynamical common core?

We claim that the answer to this is ‘yes’, on at least one plausible way of understanding what the ‘structure’ associated to a theory is. For this, recall that the Levi-Civita connection of GR is *definable* from the metric (cf. Barrett (2017)). So if one understands the ‘structure’ of a theory to include (a) all those structures to which the theory is explicitly committed—i.e. which show up between the angle brackets of its models, and (b) any structures which are definable therefrom, then the relativistic geometric trinity does indeed have a dynamical common core, which is just GR—*unlike* the non-relativistic geometric trinity (the dynamical common core of which is MG), the common core is not a distinct theory.

Several points are in order here. First one might question whether the above claim about the structural commitments of theories is correct. Second, given that  $\langle M, g_{ab}, \Phi \rangle$  is the invariant kinematical structure of the relativistic geometric trinity, one might quite reasonably ask whether it is the case that the structure of MG can be, in some sense, seen to arise as the non-relativistic limit of this. (This is especially pressing given that in §2.2 we identified NCT, rather than MG, as the non-relativistic limit of GR.) Third, if the dynamical common core of the relativistic trinity is GR, then one might wonder about the connection to Knox’s apparently similar claim that in TEGR, “the old entities from GR appear to be waiting in the wings” (Knox 2011, p. 273). We’ll now address each of these in turn.

Beginning with the first, we claim that the above characterisation of ‘structure’ (which follows the lead of Barrett (2017)) is indeed a natural and plausible one, and one which, moreover, is assumed throughout philosophical discourse on the structure of theories. To expand on what we mean here, consider the following representative statements:

- (i) A cotangent bundle  $T^*M$  comes canonically equipped with a symplectic form, so a cotangent bundle has the structure of a symplectic manifold.
- (ii) From the electromagnetic 1-form  $A_a$  on  $\langle M, \eta_{ab} \rangle$  we can define the Faraday tensor  $F_{ab}$  but not *vice versa*, therefore this 1-form has more structure than the Faraday tensor.
- (iii) Full Newtonian spacetime  $\langle M, t_a, h^{ab}, \xi^a \rangle$  has more structure than Galilean spacetime  $\langle M, t_a, h^{ab}, \nabla \rangle$ ; it has all the structure that Galilean spacetime does, but also has a standard of rest.

These statements are intuitively compelling, and moreover, strike us as *true*. But unless our claim about what one should take the structure of a theory to include is correct, it is difficult to make sense of this. E.g. on (i), nowhere in writing down ‘ $T^*M$ ’ did we explicitly write down a symplectic form  $\omega$  on  $T^*M$ ! On the other hand, our claim about the structure of theories offers an elegant explanation of *why* we should think these statements are true. E.g. on (iii), the fact that full Newtonian spacetime has all the structure that Galilean spacetime has (despite the connection not showing up explicitly in its models)



can be seen by noting that the Galilean connection is definable from the metrics and standard of rest (see Malament (2012, Proposition 4.2.4)).

Turning now to the second, consider the non-relativistic limit of  $\langle M, g_{ab}, \Phi \rangle$ . In the non-relativistic limit, the metric degenerates (see Malament (1986)), leaving us with  $\langle M, t_a, h^{ab}, \Phi \rangle$ . This structure is certainly not sufficient to formulate the dynamics of Newtonian gravitation theory—hence it is standard to take the non-relativistic limit of the formulation of GR presented in §2.1 which explicitly includes the connection, which gives us NCT. However, as noted in §3, once we have taken this limit—which gives us the geometrised Poisson equation and (7)—we still need to impose (8) for full empirical equivalence with Newtonian gravity. What happens if we also impose (8) on the spacetime  $\langle M, t_a, h^{ab}, \Phi \rangle$ ?

The basic point here is that for it even to make sense to talk of imposing (8), we still need to introduce some extra structure to  $\langle M, t_a, h^{ab}, \Phi \rangle$ . But since (8) is equivalent to the condition that (a)  $h^{ab}$  is flat, and (b) there exists a unit timelike vector field on  $M$  which is twist-free and rigid (Malament 2012, Proposition 4.2.4) this structure is strictly *less* than that of a connection—we only need the standard of rotation. So by taking the non-relativistic limit of  $\langle M, g_{ab}, \Phi \rangle$ , and then imposing the most minimal structure needed to make sense of the conditions for empirical equivalence to Newtonian gravity, one arrives naturally at the structure of Maxwellian spacetime. It is in this sense that the invariant kinematical structure of non-relativistic trinity—i.e. Maxwellian spacetime—can be seen as the non-relativistic limit of the invariant kinematical structure of the relativistic trinity  $\langle M, g_{ab}, \Phi \rangle$ .

Finally, we move to our third point—*viz.*, the connection with Knox (2011). On this, it is worth emphasising that Knox’s aims are somewhat different to ours—Knox is primarily concerned with whether GR and TEGR are in some sense *equivalent* theories, rather than with identifying their common mathematical structure. Nevertheless, there is a close relationship between Knox’s argument and our own. For example, whereas Knox takes the fact that the metric and Levi-Civita connection are still definable in TEGR (and moreover, that matter still couples to the Levi-Civita connection) to suggest that TEGR is a mere notational variant of GR, we have argued that this means that GR can be seen as the dynamical common core of the relativistic trinity—in which case, the dynamics of GR are indeed somehow ‘contained’ in the dynamics of TEGR, which is very much in the spirit of Knox’s point. Yet, *contra* Knox, and as argued by Mulder and Read (2023) and Wolf, Sanchioni, and Read (2023), there are still important reasons to regard GR, TEGR, and STEGR as distinct theories and not merely notational variants of each other—the ‘surplus’ structure that is present in TEGR and STEGR is seen by many as important for theoretical and physical reasons that lead them to be actively pursued as viable gravitational theories.

## 5 Conclusions

In this article, we’ve shown that Maxwell gravitation constitutes the mathematical common core of the recently-discovered non-relativistic geometric trinity of gravity (first presented by Wolf, Read, and Vigneron (2023)); we’ve also explained why the common core of the relativistic geometric trinity is GR, but this common core interpretation is less compelling than in the case of the non-relativistic trinity. In undertaking this work, we take ourselves to have made good on the exhortation of Lehmkuhl (2017) to explore and chart the ‘space of spacetime theories’—at least with respect to this small (albeit philosophically important!) corner of the landscape. Along the way, we have developed certain tools—e.g. the distinction between a ‘kinematical common core’ and ‘dynamical common core’ of spacetime theories—which we hope might find broader application.

There also are many future prospects; here we mention just two. First: one might be interested in whether (a) there exists an extended non-relativistic geometric trinity for the off-shell Newtonian limit presented by Hansen, Hartong, and Obers (2020) (this question was also raised by Wolf, Read, and Vigneron (2023)), and (b) if so, whether there exists a dynamical common core to this non-relativistic trinity. And second: one might wonder whether (a) there exists an *ultra*-relativistic geometric trinity obtained by taking the ultra-relativistic (i.e. roughly speaking,  $c \rightarrow 0$ ) limit of the relativistic geometric trinity, and (b) again whether there exists a dynamical common core to that trinity also.

## Acknowledgements

We are grateful to Quentin Vigneron and to Jim Weatherall for helpful discussions. E.M. acknowledges support from Balliol College, Oxford, and the Faculty of Philosophy, University of Oxford. W.W. acknowledges support from the Center for the History and Philosophy of Physics at St. Cross College, Oxford and the British Society for the Philosophy of Science. J.R. acknowledges the support of the Leverhulme Trust.

## References

- Andringa, Roel et al. (2011). “Newtonian Gravity and the Bargmann Algebra”. *Class. Quant. Grav.* 28, p. 105011. DOI: [10.1088/0264-9381/28/10/105011](https://doi.org/10.1088/0264-9381/28/10/105011). arXiv: 1011.1145 [hep-th].
- Bain, Jonathan (2006). “Spacetime Structuralism”. In: *The Ontology of Spacetime*. Ed. by Dennis Dieks. Vol. 1. Philosophy and Foundations of Physics. Elsevier, pp. 37–65. DOI: [https://doi.org/10.1016/S1871-1774\(06\)01003-5](https://doi.org/10.1016/S1871-1774(06)01003-5). URL: <https://www.sciencedirect.com/science/article/pii/S1871177406010035>.
- Barbour, Julian (2010). “Shape Dynamics: An Introduction”. In: *Proceedings of the Conference Quantum Field Theory and Gravity*. Regensburg.

- Barrett, Thomas William (2017). “What Do Symmetries Tell Us About Structure?” *Philosophy of Science* 4, pp. 617–639. DOI: 10.1086/699156.
- Capozziello, Salvatore, De Falco, Vittorio, and Ferrara, Carmen (2022). “Comparing equivalent gravities: common features and differences”. *Eur. Phys. J. C* 82.10, p. 865. DOI: 10.1140/epjc/s10052-022-10823-x. arXiv: 2208.03011 [gr-qc].
- Chen, Elliott D. (2023). *Newtonian Gravitation in Maxwell Spacetime*. URL: <http://philsci-archive.pitt.edu/22440/>.
- Dasgupta, Shamik (2016). “Symmetry as an Epistemic Notion (Twice Over)”. *British Journal for the Philosophy of Science* 67.3, pp. 837–878.
- De Haro, Sebastian and Butterfield, Jeremy (2021). “On symmetry and duality”. *Synthese* 198.4, pp. 2973–3013. DOI: 10.1007/s11229-019-02258-x. URL: <https://doi.org/10.1007/s11229-019-02258-x>.
- Dewar, Neil (2018). “Maxwell Gravitation”. *Philosophy of Science* 85, pp. 249–270.
- Duerr, Patrick and Read, James (2023). “Reconsidering Conventionalism: A Philosophy for Modern (Space-)Times?”
- Ehlers, Jürgen, Pirani, Felix AE, and Schild, Alfred (2012). “Republication of: The geometry of free fall and light propagation”. *General Relativity and Gravitation* 44.6, pp. 1587–1609.
- Fletcher, Samuel C. (2019). “On the Reduction of General Relativity to Newtonian Gravitation”. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* 68, pp. 1–15. DOI: 10.1016/j.shpsb.2019.04.005.
- Hansen, Dennis, Hartong, Jelle, and Obers, Niels A. (2019). “Action Principle for Newtonian Gravity”. *Phys. Rev. Lett.* 122 (6), p. 061106. DOI: 10.1103/PhysRevLett.122.061106. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.122.061106>.
- Hansen, Dennis, Hartong, Jelle, and Obers, Niels A. (2020). “Non-relativistic gravity and its coupling to matter”. *Journal of High Energy Physics* 2020.6, p. 145. DOI: 10.1007/JHEP06(2020)145. URL: [https://doi.org/10.1007/JHEP06\(2020\)145](https://doi.org/10.1007/JHEP06(2020)145).
- Hehl, Friedrich W. et al. (1995). “Metric-affine gauge theory of gravity: field equations, Noether identities, world spinors, and breaking of dilation invariance”. *Physics Reports* 258.1, pp. 1–171. DOI: [https://doi.org/10.1016/0370-1573\(94\)00111-F](https://doi.org/10.1016/0370-1573(94)00111-F). URL: <https://www.sciencedirect.com/science/article/pii/037015739400111F>.
- Heisenberg, Lavinia (2019). “A systematic approach to generalisations of General Relativity and their cosmological implications”. *Phys. Rept.* 796, pp. 1–113. DOI: 10.1016/j.physrep.2018.11.006. arXiv: 1807.01725 [gr-qc].
- Jacobs, Caspar (2023). “Are dynamical shifts dynamical symmetries?” *Philosophy of Science*. DOI: 10.1017/psa.2023.13.
- Jiménez, Jose Beltrán, Heisenberg, Lavinia, and Koivisto, Tomi S. (2019). “The Geometrical Trinity of Gravity”. *Universe* 5.7. ISSN: 2218-1997. DOI: 10.3390/universe5070173. URL: <https://www.mdpi.com/2218-1997/5/7/173>.

- Knox, Eleanor (2011). “Newton–Cartan Theory and Teleparallel Gravity: The Force of a Formulation”. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* 42.4, pp. 264–275. DOI: 10.1016/j.shpsb.2011.09.003.
- Knox, Eleanor (2014). “Newtonian Spacetime Structure in Light of the Equivalence Principle”. *British Journal for the Philosophy of Science* 65.4, pp. 863–880. DOI: 10.1093/bjps/axt037.
- Künzle, H. P. (May 1976). “Covariant Newtonian limit of Lorentz space-times”. *General Relativity and Gravitation* 7.5, pp. 445–457. DOI: 10.1007/BF00766139.
- Le Bihan, Baptiste and Read, James (2018). “Duality and Ontology”. *Philosophy Compass* 13, e12555.
- Lehmkuhl, Dennis (2017). “Introduction: Towards a Theory of Spacetime Theories”. In: *Towards a Theory of Spacetime Theories*. Ed. by D. Lehmkuhl, G. Schieman, and E. Scholz. Birkhäuser.
- Linnemann, Niels and Read, James (2021). “Constructive Axiomatics in Spacetime Physics Part I: Walkthrough to the Ehlers-Pirani-Schild Axiomatization”. URL: <https://arxiv.org/abs/2112.14063>.
- Malament, David (1986). “Gravity and Spatial Geometry”. In: *Studies in Logic and the Foundations of Mathematics*. Ed. by Ruth Barcan Marcus, Georg J. W. Dorn, and Paul Weingartner. Vol. 114. Elsevier, pp. 405–411. DOI: [https://doi.org/10.1016/S0049-237X\(09\)70703-7](https://doi.org/10.1016/S0049-237X(09)70703-7). URL: <https://www.sciencedirect.com/science/article/pii/S0049237X09707037>.
- Malament, David (2012). *Topics in the Foundations of General Relativity and Newtonian Gravitation Theory*. Chicago: University of Chicago Press.
- March, Eleanor (2023a). *Are Maxwell gravitation and Newton-Cartan theory theoretically equivalent?* URL: <http://philsci-archive.pitt.edu/22922/>.
- March, Eleanor (2023b). *Maxwell gravitation without reference to equivalence classes of derivative operators*. URL: <http://philsci-archive.pitt.edu/22404/>.
- Martens, Niels C. M. and Read, James (2021). “Sophistry about symmetries?” *Synthese* 199.1, pp. 315–344. DOI: 10.1007/s11229-020-02658-4. URL: <https://doi.org/10.1007/s11229-020-02658-4>.
- Matveev, Vladimir S. and Scholz, Erhard (2020). “Light cone and Weyl compatibility of conformal and projective structures”. *General Relativity and Gravitation* 52.7, p. 66. DOI: 10.1007/s10714-020-02716-9. URL: <https://doi.org/10.1007/s10714-020-02716-9>.
- Mulder, Ruward A. and Read, James (Dec. 2023). *Is spacetime curved? Assessing the underdetermination of general relativity and teleparallel gravity*. URL: <http://philsci-archive.pitt.edu/22897/>.
- Newton, Isaac (2014). *Philosophical Writings*. Ed. by Andrew Janiak. revised. Cambridge University Press.
- Oshita, Naritaka and Wu, Yi-Peng (2017). “Role of spacetime boundaries in Einstein’s other gravity”. *Phys. Rev. D* 96.4, p. 044042. DOI: 10.1103/PhysRevD.96.044042. arXiv: 1705.10436 [gr-qc].

- Pooley, Oliver (2013). “Substantialist and Relationist Approaches to Spacetime”. In: *The Oxford Handbook of Philosophy of Physics*. Ed. by Robert Batterman. Oxford University Press.
- Read, James and Teh, Nicholas J (2022). “Newtonian Equivalence Principles”. *Erkenntnis*. Forthcoming.
- Read, James and Teh, Nicholas J. (2018). “The teleparallel equivalent of Newton–Cartan gravity”. *Class. Quant. Grav.* 35.18, 18LT01. DOI: 10.1088/1361-6382/aad70d. arXiv: 1807.11779 [gr-qc].
- Rosenstock, Sarita, Barrett, Thomas William, and Weatherall, James Owen (2015). “On Einstein Algebras and Relativistic Spacetimes”. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* 52.Part B, pp. 309–316. DOI: 10.1016/j.shpsb.2015.09.003.
- Saunders, Simon (2013). “Rethinking Newton’s Principia”. *Philosophy of Science* 80, pp. 22–48.
- Schwartz, Philip K. (2023). “Teleparallel Newton–Cartan gravity”. *Class. Quant. Grav.* 40.10, p. 105008. DOI: 10.1088/1361-6382/accc02. arXiv: 2211.11796 [gr-qc].
- Teh, Nicholas J. (2018). “Recovering Recovery: On the Relationship Between Gauge Symmetry and Trautman Recovery”. *Philosophy of Science* 85.1, pp. 201–224.
- Trautman, Andrzej (1965). “Foundations and Current Problems of General Relativity”. Ed. by S. Deser and K. W. Ford.
- Wallace, David (2020). “Fundamental and Emergent Geometry in Newtonian Physics”. *British Journal for the Philosophy of Science* 71, pp. 1–32.
- Weatherall, James Owen (2016). “Maxwell-Huygens, Newton-Cartan, and Saunders-Knox Space-Times”. *Philosophy of Science* 83 (1), pp. 82–92.
- Weatherall, James Owen (2018). “A Brief Comment on Maxwell(/Newton)-Huygens] Spacetime”. *Studies in History and Philosophy of Modern Physics* 63, pp. 34–38.
- Weyl, Hermann (1921). “Zur Infinitesimalgeometrie: Einordnung der projektiven und der konformen Auffassung”. *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse* 1921, pp. 99–112.
- Wolf, William J. and Read, James (2023). “Respecting boundaries: theoretical equivalence and structure beyond dynamics”. *Eur. J. Phil. Sci.* 13.4, p. 47. DOI: 10.1007/s13194-023-00545-6. arXiv: 2302.07180 [physics.hist-ph].
- Wolf, William J., Read, James, and Teh, Nicholas J. (2023). “Edge Modes and Dressing Fields for the Newton–Cartan Quantum Hall Effect”. *Found. Phys.* 53.1, p. 3. DOI: 10.1007/s10701-022-00615-4. arXiv: 2111.08052 [cond-mat.mes-hall].
- Wolf, William J., Read, James, and Vigneron, Quentin (2023). “The Non-Relativistic Geometric Trinity of Gravity”. arXiv: 2308.07100 [gr-qc].
- Wolf, William J., Sanchioni, Marco, and Read, James (2023). “Underdetermination in Classic and Modern Tests of General Relativity”. arXiv: 2307.10074 [physics.hist-ph].